

A contribution of a $U(1)$ -reducible connection to quantum invariants of links I: R -matrix and Burau representation

L. Rozansky¹

Department of Mathematics, Yale University

10 Hillhouse Ave., P.O. Box 208283

New Haven, CT 06520-8283

E-mail address: rozansky@math.yale.edu

Abstract

We introduce a new invariant of links in S^3 which we call a contribution of a $U(1)$ -reducible connection to the colored Jones polynomial. This invariant is a formal power series in h , whose coefficients are rational functions of parameters t_j assigned to the link components. The denominators of the rational functions are the powers of the colored Alexander-Conway function. Similarly to the case of a knot, the Jones polynomial of an algebraically connected link is determined by the contribution of the $U(1)$ -reducible connection. Namely, if we substitute $t_j = (1+h)^{\alpha_j}$ in the formal power series and then expand the resulting expressions in powers of h and α_j through a special step-by-step procedure, then we obtain the expansion of the colored Jones polynomial of the link in powers of $h = q - 1$ and colors α_j . The proof of these results is based on a similarity between R -matrix and Burau representations of the braid group.

¹This work was supported by NSF Grant DMS-9704893

Contents

1	Introduction	2
1.1	Motivation	2
1.2	Notations, definitions and basic properties	5
1.2.1	Multi-index notations	5
1.2.2	Linear algebra notations	5
1.2.3	Topological notations	6
1.2.4	Alexander-Conway polynomial	8
1.2.5	Bureau representation	9
1.2.6	$SU_q(2)$ -modules and R -matrix	10
1.2.7	Colored Jones polynomial	11
1.3	Results	12
1.4	A quick recipe for calculating $U(1)$ -RC invariant	16
1.5	Path integral interpretation	18
2	Calculating the RC invariant	24
2.1	Preliminary calculations	25
2.1.1	Breaking a strand	25
2.1.2	Twisting $\check{R}$ -matrix	26
2.1.3	Parametrized and regularized trace	28
2.1.4	Verma module resolution	29
2.2	Expansion of parametrized trace in powers of h	33
2.2.1	General properties	33

2.2.2	Fixing transitions	37
2.2.3	Expansion of matrix elements	39
2.2.4	Parametrized matrices	43
2.2.5	Calculating the traces	51
2.3	Removal of parametrization	55
3	Uniqueness arguments	58
3.1	Proof of Main Theorem	58
3.2	Proof of Uniqueness Theorem	61
3.2.1	Algebraically connected links	61
3.2.2	General links	65
3.3	Proof of Corollary 3.2	67
A	Non-abelian connections in link complements	68
B	The first polynomials	73
C	The Symmetry Principle	74

1 Introduction

1.1 Motivation

The famous topological invariant of knots and links, the Alexander-Conway polynomial, can be calculated by various means. In particular, it can be calculated by Conway's skein relations or by expressing it as a determinant of a matrix related to the Burau representation of the braid group. The polynomial that comes out of these calculations is known to be

determined by the group of the knot. Therefore the Alexander-Conway polynomial has a clear topological interpretation.

The colored Jones polynomial of knots and links can be also calculated by either skein relations and cabling or by taking a trace of the product of R -matrices coming from the quantum algebra $SU_q(2)$. However, in contrast to the Alexander polynomial, the emerging expressions do not appear to have a simple structure that would point towards their topological interpretation. Therefore some tricks are needed in order to see topology inside the colored Jones polynomial.

P. Melvin and H. Morton [6] suggested to study the expansion of the colored Jones polynomial $J_\alpha(\mathcal{K}; q)$ of a knot $\mathcal{K} \subset S^3$ in powers of $q - 1$ and color α

$$J_\alpha(\mathcal{K}; q) = \sum_{n=0}^{\infty} P_{\alpha;n}(\mathcal{K}) (q - 1)^n. \quad (1.1)$$

They proved that the coefficients $P_{\alpha;n}(\mathcal{K})$ can be presented as polynomials of α and they conjectured the bound on the degree of these polynomials

$$P_{\alpha;n}(\mathcal{K}) \in \mathbb{Q}[\alpha], \quad \deg P_{\alpha;n}(\mathcal{K}) \leq n/2. \quad (1.2)$$

Since

$$q^\alpha - 1 = \alpha (q - 1) + O((q - 1)^2) \quad \text{as } q \rightarrow 1, \quad (1.3)$$

then we can rewrite the series (1.1) as

$$J_\alpha(\mathcal{K}; q) = [\alpha] \sum_{n=0}^{\infty} f_n(\mathcal{K}; q^\alpha) (q - 1)^n, \quad (1.4)$$

where

$$[\alpha] = \frac{q^{\alpha/2} - q^{-\alpha/2}}{q^{1/2} - q^{-1/2}} \quad (1.5)$$

and the coefficients $f_n(\mathcal{K}; t)$ are formal power series in $t^{1/2} - t^{-1/2}$

$$f_n(\mathcal{K}; t) = \sum_{m=0}^{\infty} d_{m;n}(\mathcal{K}) (t^{1/2} - t^{-1/2})^{2n}. \quad (1.6)$$

(only even powers of $t^{1/2} - t^{-1/2}$ participate in this expansion, because the polynomials (1.2) are known to contain only odd powers of α and $[\alpha]$ is an odd function of α).

Based on their calculations of the expansion (1.4) for torus knots, Melvin and Morton conjectured that

$$f_0(\mathcal{K}; t) = \frac{1}{\Delta_A(\mathcal{K}; t)}, \quad (1.7)$$

where $\Delta_A(\mathcal{K}; t)$ is the Alexander-Conway polynomial of $\mathcal{K}$. The conjectures of Melvin and Morton were proved by D. Bar-Natan and S. Garoufalidis [1]. We showed in [8] that the other coefficients $f_n(\mathcal{K}; t)$ of eq.(1.4) are also related to the Alexander-Conway polynomial. We proved that

$$f_n(\mathcal{K}; t) = \frac{P_n(\mathcal{K}; t)}{\Delta_A^{2n+1}(\mathcal{K}; t)}, \quad P_n(\mathcal{K}; t) \in \mathbb{Z}[t, t^{-1}]. \quad (1.8)$$

The manner in which the polynomials $P_n(\mathcal{K}; t)$ appear in the expansion (1.4), (1.8) suggests that they might present a natural extension of the Alexander-Conway polynomial and therefore may have a direct topological interpretation. Another reason to explore the polynomials $P_n(\mathcal{K}; t)$ is that the expansion

$$J_\alpha(\mathcal{K}; q) = [\alpha] \sum_{n=0}^{\infty} \frac{P_n(\mathcal{K}; q^\alpha)}{\Delta_A^{2n+1}(\mathcal{K}; q^\alpha)} (q-1)^n \quad (1.9)$$

is well-suited for studying p -adic properties of the Reshetikhin-Turaev invariant of rational homology spheres [4], [7].

The goal of this paper is to extend the expansion (1.9) to the links in S^3 . As we will see, the Melvin-Morton expansion of the colored Jones polynomial of a link can not be directly presented in the form (1.9). However, we will be able to define the analogs of polynomials $P_n(\mathcal{K}; t)$ for links whose Alexander-Conway polynomial is not identically equal to zero. We will do it in such a way that the link analog of the relation (1.9) will hold (in a certain way) if the link has sufficiently many non-zero linking numbers.

1.2 Notations, definitions and basic properties

1.2.1 Multi-index notations

For a non-negative integer A we denote $\underline{x} = (x_1, \dots, x_A)$ and for a positive k , $\underline{x}_{[k]} = (x_1, \dots, x_{A-k})$ and $\underline{x}_{(k)} = (x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_A)$. We also use the following notations

$$\begin{aligned} y\underline{x} &= (yx_1, \dots, yx_A), & \underline{x}^y &= (x_1^y, \dots, x_A^y), & \underline{x}\underline{y} &= (x_1y_1, \dots, x_Ay_A), \\ \{f(\underline{x})\} &= \prod_{j=1}^A f(x_j), & |\underline{x}| &= \sum_{j=1}^A x_j, & \underline{x}^{\underline{y}} &= \prod_{j=1}^A x_j^{y_j}. \end{aligned} \quad (1.10)$$

For a constant C we denote

$$\underline{C} = (C, \dots, C). \quad (1.11)$$

We assume concatenation of multi-indices

$$x, \underline{y} = (x, y_1, \dots, y_A), \quad \underline{x}, \underline{y} = (x_1, \dots, x_A, y_1, \dots, y_B) \quad \text{if } \underline{y} = (y_1, \dots, y_B). \quad (1.12)$$

Also $\underline{x} = x$ means $x_j = x$ for all $1 \leq j \leq A$.

Let $f(\underline{x})$ be a meromorphic function of the variables $\underline{x}$ which may have singularities in the vicinity of $\underline{x} = 0$. $[f(\underline{x})]_{(\underline{x})}$ denotes a result of sequential expansion of $f(\underline{x})$ in powers of $\underline{x}$ when we start with expanding in x_L and finish with expanding in x_1 .

1.2.2 Linear algebra notations

Let V be a linear space with a basis f_m , $m \geq 0$. If an operator O acts on V , then we denote its matrix elements as O_m^n so that

$$O(f_m) = \sum_n O_m^n f_n. \quad (1.13)$$

Suppose that V_1 and V_2 are two linear spaces with bases f_m , $m \geq 0$. Let O be an operator which maps the tensor product $V_1 \otimes V_2$ either into itself or into $V_2 \otimes V_1$. We denote the matrix elements of O as $O_{m_1, m_2}^{n_1, n_2}$ so that

$$O(f_{m_1} \otimes f_{m_2}) = \sum_{n_1, n_2} O_{m_1, m_2}^{n_1, n_2} f_{n_1} \otimes f_{n_2}. \quad (1.14)$$

If V is a linear space, then for a positive integer N we denote

$$V^{\otimes N} = \underbrace{V \otimes \cdots \otimes V}_{N \text{ times}}. \quad (1.15)$$

Similarly, if O is an operator acting on V , then $O^{\otimes N}$ denotes a corresponding operator acting on $V^{\otimes N}$.

Let O be an operator acting on a tensor product of linear spaces $V \otimes W$. If we take a ‘partial’ trace of O over the space W , then we obtain an operator acting on V . We denote it as $\text{Tr}_W O$. Suppose that there is a subspace $U \subset V$ such that $(\text{Tr}_W O)(U) \subset U$. This means that the action of $\text{Tr}_W O$ on V can be restricted to U . We denote the trace of $\text{Tr}_W O$ over U simply as $\text{Tr}_{U \otimes W} O$:

$$\text{Tr}_{U \otimes W} O = \text{Tr}_U (\text{Tr}_W O) \quad (1.16)$$

Remark 1.1 Note that this notation is a bit misleading because, generally speaking, the action of O on $U \otimes W$ is not well-defined. However, we hope that there will be no confusion in the context of our calculations.

1.2.3 Topological notations

In this paper $\mathcal{L}$ (or $\mathcal{L}'$) usually denotes an L (or L')-component link in S^3 , while $\mathcal{L}_j$, $1 \leq j \leq L$ (or $\mathcal{L}'_j$, $1 \leq j \leq L'$) denote its components. The numbers l_{ij} , $1 \leq i, j \leq L$ (or l'_{ij} , $1 \leq i, j \leq L'$) are the linking numbers of $\mathcal{L}$ (or $\mathcal{L}'$). The numbers l''_{ij} , denote the linking numbers between $\mathcal{L}_i \in \mathcal{L}$ and $\mathcal{L}'_j \in \mathcal{L}'$. We denote

$$\text{lk}(\mathcal{L}; \underline{x}) = \frac{1}{2} \sum_{1 \leq i < j \leq L} l_{ij} x_i x_j \quad (1.17)$$

For $1 \leq k \leq L - 1$, $\mathcal{L}_{[k]}$ denotes a sublink of $\mathcal{L}$ which consists of components $\mathcal{L}_j$, $1 \leq j \leq L - k$ and for $1 \leq k \leq L$, $\mathcal{L}_{(k)}$ denotes the link $\mathcal{L}$ with the component $\mathcal{L}_k$ removed.

Definition 1.2 A link $\mathcal{L}$ is *algebraically connected* to another link $\mathcal{L}'$ if either one of the links is empty, or at least one of the linking numbers l''_{ij} is non-zero.

Definition 1.3 A link $\mathcal{L}$ is called algebraically connected if it can not be presented as a union of two links which are not algebraically connected to each other.

Definition 1.4 The indexing of the components of an algebraically connected link $\mathcal{L}$ is called *admissible* if $\mathcal{L}_{[k]}$ is algebraically connected for any $1 \leq k \leq L - 1$.

Lemma 1.5 If $\mathcal{L}$ is an algebraically connected L -component link, then for any selected component of $\mathcal{L}$ there exists an admissible indexing such that the selected component becomes $\mathcal{L}_1$.

Proof. We leave the proof to the reader. □

Let $\mathcal{B}$ be an N -component braid. Its *closure* is constructed by connecting the upper ends of its strands with the lower ends in the same order. For any link $\mathcal{L}$ there exists a braid $\mathcal{B}_{\mathcal{L}}$ whose closure is $\mathcal{L}$. The numbers $\underline{N} = (N_1, \dots, N_L)$ denote the numbers of strands of $\mathcal{B}$ which belong to the link components $\mathcal{L}_j$ of $\mathcal{L}$.

In this paper we will always assume that the strands of $\mathcal{L}$ appear at the bottom and top of $\mathcal{B}$ in a particular order: if $i < j$, then the strands of $\mathcal{L}_i$ precede the strands of $\mathcal{L}_j$. If we present a union of two links $\mathcal{L} \cup \mathcal{L}'$ as a closure of a braid $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$, then we also assume that the strands of $\mathcal{L}$ precede the strands of $\mathcal{L}'$. $\underline{N}' = (N_1, \dots, N_{L'})$ denote the numbers of strands in the link components of $\mathcal{L}'$.

For any link $\mathcal{L}$ one can always find a braid $\mathcal{B}_{\mathcal{L}}$ which satisfies the ordering property. Note, that this property of $\mathcal{B}_{\mathcal{L}}$ is completely unnecessary for the calculations that we will perform in this paper, but it will simplify our notations.

We say that an elementary crossings of $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ is of $(\mathcal{L}, \mathcal{L})$, $(\mathcal{L}, \mathcal{L}')$, $(\mathcal{L}', \mathcal{L})$ or $(\mathcal{L}', \mathcal{L}')$ type depending on whether the incoming strands belong to $\mathcal{L}$ or $\mathcal{L}'$.

1.2.4 Alexander-Conway polynomial

Let $\mathcal{K}$ be a knot in S^3 . $\Delta_A(\mathcal{K}; t)$ denotes its Alexander-Conway polynomial. This polynomial satisfies the skein relation of Fig. 1 and it is normalized by the condition

$$\Delta_A(\mathcal{K}; 1) = 1. \quad (1.18)$$

$$\Delta_A \left(\text{positive crossing} ; t \right) - \Delta_A \left(\text{negative crossing} ; t \right) = (t^{1/2} - t^{-1/2}) \Delta_A \left(\text{two parallel strands} ; t \right).$$

Figure 1: The skein relation for the Alexander-Conway polynomial

For an *oriented* L -component link $\mathcal{L} \subset S^3$ we define the Alexander-Conway function $\nabla_A(\mathcal{L}; \underline{t})$, $\underline{t} = (t_1, \dots, t_L)$ according to the definition of [9] except that our $\underline{t}$ is equal to $\underline{t}^2$ of Turaev. The function $\nabla_A(\mathcal{L}; \underline{t})$ is almost a polynomial:

$$\nabla_A(\mathcal{L}; \underline{t}) \in \begin{cases} \frac{\mathbb{Z}[t_1, t_1^{-1}]}{t_1^{1/2} - t_1^{-1/2}} & \text{if } L = 1 \\ \mathbb{Z}[\underline{t}^{1/2}, \underline{t}^{-1/2}] & \text{if } L \geq 2. \end{cases} \quad (1.19)$$

It satisfies the following properties:

$$\nabla_A(\mathcal{K}; t) = \frac{\Delta_A(\mathcal{K}; t)}{t^{1/2} - t^{-1/2}} \quad \text{for } \mathcal{K} \text{ being a knot } \mathcal{K}, \quad (1.20)$$

$$\nabla_A(\text{Hopf link}; t_1, t_2) = 1, \quad (1.21)$$

$$\nabla_A(\mathcal{L}; \underline{t}^{-1}) = (-1)^L \nabla_A(\mathcal{L}; \underline{t}). \quad (1.22)$$

If we change the orientation of a component $\mathcal{L}_j$ and denote the new oriented link as $\mathcal{L}'$, then

$$\nabla_A(\mathcal{L}'; \underline{t}) = -\nabla_A(\mathcal{L}; t_1, \dots, t_j^{-1}, \dots, t_L). \quad (1.23)$$

The Alexander-Conway function satisfies the Torres formula: if $L \geq 2$, then for i , $1 \leq i \leq L$

$$\nabla_A(\mathcal{L}; \underline{t})|_{t_i=1} = \left(\prod_{\substack{1 \leq j \leq L \\ j \neq i}} t_j^{l_{ij}/2} - \prod_{\substack{1 \leq j \leq L \\ j \neq i}} t_j^{-l_{ij}/2} \right) \nabla_A(\mathcal{L}_{(i)}; \underline{t}_{(i)}). \quad (1.24)$$

A correction factor

$$\Phi_A(\mathcal{L}; \underline{t}) = \prod_{j=1}^L t_j^{-\frac{1}{2}(\sum_{i: i \neq j} l_{ij} + 1)} \quad (1.25)$$

allows us to remove the fractional powers of $\underline{t}$ from $\nabla_A(\mathcal{L}; \underline{t})$: if $L \geq 2$ then

$$\Phi_A(\mathcal{L}; \underline{t}) \nabla_A(\mathcal{L}; \underline{t}) \in \mathbb{Z}[\underline{t}, \underline{t}^{-1}]. \quad (1.26)$$

1.2.5 Burau representation

The Alexander-Conway function of a link can be calculated with the help of Burau representation of the braid group. Let $\mathcal{B}_{\mathcal{L}}$ be an N -strand braid whose closure is a link $\mathcal{L}$. Consider a linear space $\mathbb{C}^N$ whose basis vectors $\underline{e} = (e_1, \dots, e_N)$ are associated with N positions of the braid strands. To a positive (negative) elementary braid in positions $j, j+1$ we associate an operator $\rho_+[t_1, t_2]$ ($\rho_-[t_1, t_2]$) acting on the vectors e_j, e_{j+1}

$$\rho_+[t_1, t_2] = \begin{pmatrix} 1 - t_2^{-1} & t_1^{-1} \\ 1 & 0 \end{pmatrix}, \quad \rho_-[t_1, t_2] = \begin{pmatrix} 0 & 1 \\ t_2 & t_1^{-1}t_2(1 - t_1) \end{pmatrix}, \quad (1.27)$$

where t_1 and t_2 denote the variables related to the incoming link components. Note that $\rho_+^{-1}[t_1, t_2] = \rho_-[t_2, t_1]$. If we present the braid $\mathcal{B}_{\mathcal{L}}$ as a product of elementary braids and then replace those braids with operators $\rho_{\pm}$, then we obtain an operator $B_{\mathcal{L}}[\underline{t}]$ which acts on $\mathbb{C}^N$. Let Q be an operator which acts on $\mathbb{C}^N$ as

$$Q(e_1) = 0, \quad Q(e_j) = e_j \quad \text{for } j \neq 1. \quad (1.28)$$

Then

$$\nabla_A(\mathcal{L}; \underline{t}) = \Phi_1(\mathcal{B}_{\mathcal{L}}; \underline{t}) \frac{\det_{\mathbb{C}^N}(1 - QB_{\mathcal{L}}[\underline{t}])}{t_1^{1/2} - t_1^{-1/2}}, \quad (1.29)$$

where

$$\Phi_1(\mathcal{B}_{\mathcal{L}}; \underline{t}) = t_1^{1/2} \prod_{j=1}^L t_j^{\frac{1}{2}(-N_j + \sum_{i=1}^L l_{ij})}. \quad (1.30)$$

1.2.6 $SU_q(2)$ -modules and R -matrix

The quantum invariants in 3-dimensional topology depend on a ‘quantum deformation parameter’ q . We do not assume that q is a root of unity unless stated explicitly. We also use a parameter

$$h = q - 1. \quad (1.31)$$

Let X , Y and H be the generators of the quantum group $SU_q(2)$ which satisfy the commutation relations

$$[H, X] = 2X, \quad [H, Y] = -2Y, \quad [X, Y] = [H]. \quad (1.32)$$

In the last equation of (1.32) we used a standard notation

$$[x] = \frac{q^{x/2} - q^{-x/2}}{q^{1/2} - q^{-1/2}}. \quad (1.33)$$

We choose a basis f_m , $0 \leq m \leq \gamma - 1$ of a γ -dimensional $SU_q(2)$ module V_γ in such a way that

$$H f_m = (\gamma - 2m - 1) f_m, \quad (1.34)$$

$$X f_m = [m] f_{m-1}, \quad (1.35)$$

$$Y f_m = [\gamma - m - 1] f_{m+1}, \quad (1.36)$$

We will also use an auxiliary operator $\hat{m}$

$$\hat{m}(f_m) = m f_m. \quad (1.37)$$

R -matrix acts on a tensor product of two $SU_q(2)$ modules $V_{\gamma_1} \otimes V_{\gamma_2}$

$$R(f_{m_1} \otimes f_{m_2}) = \sum_{0 \leq n \leq \max\{m_1, \gamma_2 - m_2 - 1\}} R_{m_1, m_2}^{m_1 - n, m_2 + n} f_{m_1 - n} \otimes f_{m_2 + n}, \quad (1.38)$$

where

$$\begin{aligned} R_{m_1, m_2}^{m_1 - n, m_2 + n} &= (q^{1/2} - q^{-1/2})^n \frac{[\gamma_2 - m_2 - 1]!}{[\gamma_2 - m_2 - 1 - n]!} \frac{[m_1]!}{[m_1 - n]! [n]!} \\ &\quad \times q^{\frac{1}{4}((\gamma_1 - 2m_1 - 1)(\gamma_2 - 2m_2 - 1) - n(\gamma_1 - \gamma_2 - 2m_1 + 2m_2 + n + 1))}. \end{aligned} \quad (1.39)$$

1.2.7 Colored Jones polynomial

Let $\mathcal{L} \subset S^3$ be an L -component link which is colored by positive integers $\underline{\alpha} = (\alpha_1, \dots, \alpha_L)$. We present $\mathcal{L}$ as a closure of an N -strand braid $\mathcal{B}_{\mathcal{L}}$. We index the strands of $\mathcal{B}_{\mathcal{L}}$ according to the order in which they appear at the bottom of the braid.

Consider a tensor product of $SU_q(2)$ modules ‘attached’ to the braid strands

$$V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha}} = \bigotimes_{j=1}^L V_{\alpha_j}^{\otimes N_j}. \quad (1.40)$$

We can construct an action of the braid $\mathcal{B}$ on $V_{\mathcal{B};\underline{\alpha}}$ if we present $\mathcal{B}$ as a product of positive and negative elementary braids and replace each positive braid with the operator

$$\check{R} = P R, \quad P: V_{\gamma_1} \otimes V_{\gamma_2} \rightarrow V_{\gamma_2} \otimes V_{\gamma_1}, \quad P(f_{m_1} \otimes f_{m_2}) = f_{m_2} \otimes f_{m_1} \quad (1.41)$$

acting on a product of $SU_q(2)$ modules assigned to the braided pair of strands, while replacing each negative elementary braid with the inverse operator $\check{R}^{-1}$. We denote the resulting operator as

$$B_{\mathcal{L}}: V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha}} \rightarrow V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha}}. \quad (1.42)$$

We introduce the ‘quantum trace’ operator

$$Q_q = \left(q^{H/2}\right)^{\otimes N}. \quad (1.43)$$

The colored Jones polynomial of $\mathcal{L}$ is defined as a trace over $V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha}}$

$$J_{\underline{\alpha}}(\mathcal{L}; q) = q^{-\phi_{\text{sl}}} \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha}}} Q_q B_{\mathcal{L}}. \quad (1.44)$$

The factor $q^{-\phi_{\text{sl}}}$ is necessary to correct self-linking of the components of $\mathcal{L}$

$$\phi_{\text{sl}}(\mathcal{L}) = \frac{1}{4} \sum_{j=1}^L l_{jj} (\alpha_j^2 - 1), \quad (1.45)$$

where l_{jj} is the number of positive self-crossings minus the number of negative self-crossings of the strands of $\mathcal{B}$ which constitute $\mathcal{L}_j$.

Being a polynomial of $q^{1/2}, q^{-1/2}$, the colored Jones polynomial of $\mathcal{L}$ can be expanded in powers of $h = q - 1$. Melvin and Morton [6] proved that the coefficients in this expansion are polynomials of the color variables

$$J_{\underline{\alpha}}(\mathcal{L}; q) = \sum_{n=0}^{\infty} P_{\underline{\alpha};n}(\mathcal{L}) h^n, \quad (1.46)$$

$$P_{\underline{\alpha};n}(\mathcal{L}) \in \mathbb{Q}[\underline{\alpha}], \quad \deg_{\underline{\alpha}} P_{\underline{\alpha};n}(\mathcal{L}) \leq 2n + 1.$$

and the polynomials $P_{\underline{\alpha};n}(\mathcal{L})$ are odd functions of all variables $\underline{\alpha}$. We call eq. (1.46) *the Melvin-Morton expansion*.

1.3 Results

The purpose of this paper is to prove the following

Main Theorem *Let $\mathcal{L}$ be an oriented L -component link in S^3 , whose Alexander-Conway function is not identically equal to zero*

$$\nabla_A(\mathcal{L}; \underline{t}) \not\equiv 0, \quad \underline{t} = (t_1, \dots, t_L). \quad (1.47)$$

Let $\mathcal{L}'$ be an L' -component link in S^3 with positive integers $\underline{\beta} = (\beta_1, \dots, \beta_{L'})$ assigned to its components. Then there exists a unique sequence of polynomial invariants

$$P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}) \in \mathbb{Z}[\underline{t}^{1/2}, \underline{t}^{-1/2}, 1/2], \quad n \geq 0, \quad (1.48)$$

such that an expansion of $P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})$ in powers of $\underline{t} - 1$ has the form

$$P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}) = \sum_{\underline{m} \geq 0} p_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) (\underline{t} - 1)^{\underline{m}}, \quad p_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \in \mathbb{Q}[\underline{\beta}], \quad (1.49)$$

and the formal power series in h

$$\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h) = \begin{cases} \sum_{n=0}^{\infty} P_{\underline{\beta};n}(\mathcal{L}') h^n & \text{if } L = 0, \\ \frac{1}{h} \frac{1}{\nabla_A(\mathcal{L}; t_1)} \sum_{n=0}^{\infty} \frac{P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; t_1)}{\Delta_A^{2n}(\mathcal{L}; t_1)} h^n & \text{if } L = 1, \\ \frac{1}{h} \frac{1}{\nabla_A(\mathcal{L}; \underline{t})} \sum_{n=0}^{\infty} \frac{P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})}{\nabla_A^{2n}(\mathcal{L}; \underline{t})} h^n & \text{if } L \geq 2 \end{cases} \quad (1.50)$$

satisfies the following properties:

(1) If the link $\mathcal{L}$ is empty (that is, if $L = 0$), then $\check{J}_{\underline{\beta}}^{(r)}(\emptyset, \mathcal{L}'; h)$ is the Melvin-Morton expansion (1.46) of $J_{\underline{\beta}}(\mathcal{L}'; q)$

$$\check{J}_{\underline{\beta}}^{(r)}(\emptyset, \mathcal{L}'; h) = J_{\underline{\beta}}(\mathcal{L}'; q) \Big|_{q=1+h} \quad \text{in } \mathbb{Q}[\underline{\beta}][[h]]. \quad (1.51)$$

(2) For any j , $1 \leq j \leq L'$,

$$\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h) \Big|_{\beta_j=1} = \check{J}_{\underline{\beta}_{(j)}}^{(r)}(\mathcal{L}, \mathcal{L}'_{(j)}; \underline{t}; h). \quad (1.52)$$

(3) Let $\mathcal{L}$ be a non-empty link. Suppose that for a number i , $1 \leq i \leq L$, the link component $\mathcal{L}_i$ is algebraically connected to $\mathcal{L}_{(i)}$ and $\nabla_A(\mathcal{L}_{(i)}; \underline{t}_{(i)}) \neq 0$. Then

$$\check{J}_{\beta_0, \underline{\beta}}^{(r)}(\mathcal{L}_{(i)}, \mathcal{L}_i \cup \mathcal{L}'; \underline{t}_{(i)}; h) = \begin{cases} \sum_{\mu=\pm 1} \mu \left(\prod_{\substack{1 \leq j \leq L \\ j \neq i}} t_j^{l_{ij}} \right)^{\mu\beta_0/2} \check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h) \Big|_{t_i=q^{\mu\beta_0}} & \text{if } L \geq 2, \\ \check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q^{\beta_0}; h) & \text{if } L = 1. \end{cases} \quad (1.53)$$

Thus defined, the polynomials $P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})$ have additional properties:

$$P_{\underline{\beta};0}(\mathcal{L}, \mathcal{L}'; \underline{t}) = \begin{cases} \{\underline{\beta}\} & \text{if } L = 0, \\ \prod_{j=1}^{L'} \frac{\left(\prod_{i=1}^L t_i^{l_{ij}} \right)^{\beta_j/2} - \left(\prod_{i=1}^L t_i^{l_{ij}} \right)^{-\beta_j/2}}{\left(\prod_{i=1}^L t_i^{l_{ij}} \right)^{1/2} - \left(\prod_{i=1}^L t_i^{l_{ij}} \right)^{-1/2}} & \text{if } L \geq 1, \end{cases} \quad (1.54)$$

$$P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}^{-1}) = P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}), \quad (1.55)$$

and if we reverse the orientation of a link component $\mathcal{L}_j$, then for the new oriented link $\bar{\mathcal{L}}$

$$P_{\underline{\beta};n}(\bar{\mathcal{L}}, \mathcal{L}'; \underline{t}) = P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; t_1, \dots, t_j^{-1}, \dots, t_L). \quad (1.56)$$

The polynomials $p_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta})$ contain only odd powers of β_j

$$p_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \beta_1, \dots, -\beta_j, \dots, \beta_L) = -p_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta}), \quad (1.57)$$

and their degree is bounded

$$\deg p_{\underline{m};n} \leq \begin{cases} |\underline{m}| + 2n + 1 & \text{if } L = 1, \\ |\underline{m}| + 4n + 1 & \text{if } L \geq 1. \end{cases} \quad (1.58)$$

Finally, if we select an orientation for the components of $\mathcal{L}'$, then the formal power series (1.50) can be rewritten as

$$\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h) = \begin{cases} q^{\phi_{\text{lk}}(\mathcal{L}, \mathcal{L}'; \underline{\beta})} \sum_{n=0}^{\infty} P'_{\underline{\beta}; n}(\mathcal{L}') h^n & \text{if } L = 0, \\ \frac{1}{h} q^{\phi_{\text{lk}}(\mathcal{L}, \mathcal{L}'; \underline{\beta})} \frac{\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; t_1)}{\nabla_A(\mathcal{L}; t_1)} \sum_{n=0}^{\infty} \frac{P'_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; t_1)}{\Delta_A^{2n}(\mathcal{L}; t_1)} h^n & \text{if } L = 1, \\ \frac{1}{h} q^{\phi_{\text{lk}}(\mathcal{L}, \mathcal{L}'; \underline{\beta})} \frac{\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{t})}{\nabla_A(\mathcal{L}; \underline{t})} \sum_{n=0}^{\infty} \frac{P'_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{t})}{\nabla_A^{2n}(\mathcal{L}; \underline{t})} h^n & \text{if } L \geq 2, \end{cases} \quad (1.59)$$

where

$$\begin{aligned} & \phi_{\text{lk}}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \\ &= \frac{1}{2} \left(L + \sum_{1 \leq i < j \leq L} l_{ij} + \sum_{1 \leq i < j \leq L'} l'_{ij} (\beta_i - 1)(\beta_j - 1) - \sum_{\substack{1 \leq i \leq L \\ 1 \leq j \leq L'}} l''_{ij} (\beta_j - 1) - \sum_{j=1}^{L'} (\beta_j - 1) \right) \end{aligned} \quad (1.60)$$

$$\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{t}) = \prod_{i=1}^L t_i^{\frac{1}{2} \sum_{j=1}^{L'} l''_{ij} (\beta_j - 1)}, \quad (1.61)$$

and

$$P'_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{t}) \in \mathbb{Z}[\underline{t}, \underline{t}^{-1}] \quad (1.62)$$

for any integer $\underline{\beta}$.

Remark 1.6 A combination of eqs. (1.55) and (1.56) implies that if we take a link $\mathcal{L}$ and reverse the orientation of all its components, then for the resulting link $\bar{\mathcal{L}}$

$$P_{\underline{\beta}; n}(\bar{\mathcal{L}}, \mathcal{L}'; \underline{t}) = P_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{t}). \quad (1.63)$$

The uniqueness of the polynomials (1.48) means that the formal power series (1.50) is an invariant of a pair of (oriented and unoriented) links $\mathcal{L}, \mathcal{L}'$.

Definition 1.7 We call the expression

$$J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q) = q^{\text{lk}(\mathcal{L}; \underline{\alpha})} \check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q^{\underline{\alpha}}; h) \quad (1.64)$$

the $U(1)$ reducible connection contribution to the Jones polynomial, or simply the $U(1)$ -RC invariant.

Corollary 1.8 *Suppose that the link $\mathcal{L}$ is algebraically connected and that it has admissible indexing. Let us expand the sum*

$$\sum_{\substack{\mu_j = \pm 1 \\ 2 \leq j \leq L \\ \mu_1 = 1}} \{\underline{\mu}\} J_{\underline{\mu}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q) \quad (1.65)$$

in powers of h and $\underline{\alpha}$ going in sequence of L steps, at $(L-i+1)$ -st step ($1 \leq i \leq L$) expanding in powers of α_i and h by using the formulas

$$q^{\alpha_i} = (1+h)^{\alpha_i} = \sum_{n=0}^{\infty} \binom{\alpha_i}{n} h^n, \quad (1.66)$$

$$q^{\alpha_i \alpha_j} = (1+(q^{\alpha_j} - 1))^{\alpha_i} = \sum_{n=0}^{\infty} \binom{\alpha_i}{n} (q^{\alpha_j} - 1)^n \quad \text{for } 1 \leq j < i. \quad (1.67)$$

The resulting power series coincides with the Melvin-Morton expansion of the colored Jones polynomial $J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q)$

$$\left[\sum_{\substack{\mu_j = \pm 1 \\ 2 \leq j \leq L \\ \mu_1 = 1}} \{\underline{\mu}\} J_{\underline{\mu}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q) \right]_{(\underline{\alpha})} = J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q) \quad \text{in } \mathbb{Q}[\underline{\alpha}, \underline{\beta}][[h]]. \quad (1.68)$$

Remark 1.9 Since the Alexander-Conway function $\nabla_A(\mathcal{L} \cup \mathcal{L}'; \underline{t})$ may have zeroes in the vicinity of $\underline{t} = 1$, then the order of expanding $J_{\underline{\mu}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q)$ in powers of $\underline{\alpha}$ is important. In fact, the order suggested in Corollary 1.8 guarantees that no negative powers of $\underline{\alpha}$ appear at any stage in the expansion of the sum over $\underline{\mu}$.

Proof of Corollary 1.8. Since $\mathcal{L}$ is algebraically connected, then for any j , $2 \leq j \leq L$, the link component $\mathcal{L}_j$ has non-zero linking number with at least one of the components of $\mathcal{L}_{[j]}$. Therefore we can apply Claim 3 of Main Theorem at each step of the expansion of the sum (1.65). After we complete the expansion in powers of $\underline{\alpha}$, we apply Claim 2 which leads us to the claim of Corollary 1.8. $\square$

Remark 1.10 Suppose that $q = e^{2\pi i/K}$, where K is a positive integer and the colors $\underline{\alpha}$ are positive integers between 1 and $K-1$. Then Equations (1.58) and (1.68) lead to a

simple proof of the so-called Symmetry Principle established by R. Kirby and P. Melvin [5], which describes how the colored Jones polynomial of a link changes when one of the colors is changed from α_j to $K - \alpha_j$. We give a precise formulation and a detailed proof of the Symmetry Principle in Appendix C.

1.4 A quick recipe for calculating $U(1)$ -RC invariant

Our proof of Main Theorem is constructive. In Section 2 we give a detailed recipe for calculating the $U(1)$ -RC invariant $\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h)$. The presence of the link $\mathcal{L}'$ makes this calculation more involved, so here we will explain briefly how to obtain the $U(1)$ -RC invariant $\check{J}^{(r)}(\mathcal{L}, \emptyset; \underline{t}; h)$ in case when $\mathcal{L}'$ is empty.

First of all, we have to get the polynomials $T_{j,k}^{(\pm)}(m_1, m_2, n)$, described in Theorem 2.11. These polynomials come from expanding the following expressions

$$\begin{aligned} X^{(+)}(t_1, t_2; m_1, m_2, n; q) \\ = q^{(m_1+1)m_2+\frac{1}{2}n(n+1)} \prod_{l=1}^n \left(\frac{(q^{-m_2-l} - t_2^{-1})(q^{m_1-n+l} - 1)}{q^l - 1} \right), \end{aligned} \quad (1.69)$$

$$\begin{aligned} X^{(-)}(t_1, t_2; m_1, m_2, n; q) \\ = q^{-m_1(m_2+1)-\frac{1}{2}n(n+1)} \prod_{l=1}^n \left(\frac{(q^{m_1+l} - t_1)(q^{-m_2+n-l} - 1)}{q^{-l} - 1} \right) \end{aligned} \quad (1.70)$$

in powers of $h = q - 1$ (while keeping all other variables fixed):

$$\begin{aligned} X^{(+)}(t_1, t_2; m_1, m_2, n; q) \\ = \binom{m_1}{n} (1 - t_2^{-1})^n \sum_{j=0}^n \left(\frac{\prod_{l=0}^{j-1} (n-l)}{(1 - t_2^{-1})^j} \sum_{k=0}^{\infty} T_{j,k}^{(+)}(m_1, m_2, n) h^{j+k} \right), \end{aligned} \quad (1.71)$$

$$\begin{aligned} X^{(-)}(t_1, t_2; m_1, m_2, n; q) \\ = \binom{m_2}{n} (1 - t_1)^n \sum_{j=0}^n \left(\frac{\prod_{l=0}^{j-1} (n-l)}{(1 - t_1)^j} \sum_{k=0}^{\infty} T_{j,k}^{(-)}(m_1, m_2, n) h^{j+k} \right). \end{aligned} \quad (1.72)$$

We will also need the coefficients $C_{k,n}$ which appear in the expansion

$$\frac{1}{n!} ((1+h)^{-1/2} - 1)^n = \sum_{k \geq n} C_{k,n} h^k. \quad (1.73)$$

Suppose that an L -component link $\mathcal{L}$ can be constructed as a closure of an N -strand braid $\mathcal{B}_{\mathcal{L}}$. Let us present $\mathcal{B}_{\mathcal{L}}$ as a product of N' elementary braids

$$\mathcal{B}_{\mathcal{L}} = \prod_{l=1}^{N'} \sigma_{p(l)}^{(s(l))}, \quad (1.74)$$

where $\sigma_p^{(\pm)}$ is a positive (negative) elementary braid which twists the strands in the p -th and $(p+1)$ -th positions, while $p(l)$ and $s(l)$ are the functions describing the positions and signs of the elementary braids which comprise a presentation for $\mathcal{B}_{\mathcal{L}}$. We are going to map this presentation into an operator acting on an N -dimensional space $\mathbb{C}^N$ with a chosen basis $\underline{e} = (e_1, \dots, e_N)$. Let $\hat{R}_p^{\pm}[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4]$, $1 \leq p \leq N-1$ be the matrices (2.160), (2.161) acting on the basis elements e_p, e_{p+1} . Then we define the operator

$$\hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}] = \prod_{l=1}^{N'} \hat{R}_{p(l)}^{s(l)}[t_{n_1(l)}, t_{n_2(l)}, \epsilon_{1,l}, \epsilon_{2,l}, \epsilon_{3,l}, \epsilon_{4,l}], \quad (1.75)$$

where $\underline{\epsilon} = (\epsilon_{j,l} \mid 1 \leq j \leq 4, 1 \leq l \leq N')$ and the functions $n_1(l), n_2(l)$ indicate to which link components belong the strands which come into the l -th elementary braid $\sigma_{p(l)}^{(s(l))}$. We also need an operator

$$\hat{Q} = \text{diag}(0, 1, \dots, 1), \quad (1.76)$$

which projects out e_1 .

Next, we construct a formal power series in h whose coefficients are polynomials of the variables λ and $\underline{\epsilon}$

$$D(\lambda, \underline{\epsilon}; h) = \left(\sum_{\substack{k \geq 0 \\ 0 \leq n \leq k}} C_{k,n} \lambda^n h^k \right) \prod_{l=1}^{N'} \left(\sum_{j,k=0}^{\infty} T_{j,k}^{(s(l))}(\epsilon_{1,l}, \epsilon_{2,l}, \epsilon_{3,l}) \epsilon_{4,l}^j h^{j+k} \right). \quad (1.77)$$

Now

$$\begin{aligned} \check{J}^{(r)}(\mathcal{L}, \emptyset; \underline{t}; h) &= \frac{1 - t_1^{-1}}{1 - q^{-1}} q^{\frac{1}{2} \sum_{j=1}^L (l_{jj} - N_j)} \prod_{j=1}^L t_j^{\frac{1}{2} (N_j - \sum_{i=1}^L l_{ij})} \\ &\quad \times D(\lambda \partial_{\lambda}, \partial_{\underline{\epsilon}}; h) \frac{1}{\det_{\mathbb{C}^N} (1 - \lambda \hat{Q} \hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}])} \Bigg|_{\substack{\underline{\epsilon}=0 \\ \lambda=1}}, \end{aligned} \quad (1.78)$$

where N_j is the number of the strands of the braid $\mathcal{B}_{\mathcal{L}}$ which are part of the link component $\mathcal{L}_j$ and l_{jj} is the self-linking number of $\mathcal{L}_j$ defined by the closure of $\mathcal{B}_{\mathcal{L}}$ (in other words, l_{jj} equals the number of positive self-crossings of the strands of $\mathcal{L}_j$ minus the number of negative self-crossings).

Remark 1.11 The formula (1.78) and the detailed calculations of subsections 2.2.3 – 2.2.5 may seem rather ‘unnatural’. The natural way of calculating the $U(1)$ -RC invariant through R -matrix formula for the colored Jones polynomial (1.44) is to use the machinery of free field representation for $SU_q(2)$ generators and the perturbative expansion of quantum field theory. In this paper we tried to avoid any explicit reference to these methods and stick to simple algebra. However, we hope to explain the application of QFT methods in a future publication dealing with the colored Jones polynomials associated to simple Lie algebras of higher rank.

1.5 Path integral interpretation

E. Witten showed [10] that the colored Jones polynomial of a link S^3 can be expressed in terms of a path integral over $SU(2)$ connections. For a positive integer K and an L -component link $\mathcal{L}$ in a 3-manifold M consider the following path integral which goes over connections A in the trivial $SU(2)$ bundle over M

$$Z_{\underline{\alpha}}(M, \mathcal{L}; K) = \int \mathcal{D}A \, e^{(iK/2\pi)S_{\text{CS}}[A]} \prod_{j=1}^L \text{Tr}_{\alpha_j} \text{Pexp} \left(\oint_{\mathcal{L}_j} A dx \right). \quad (1.79)$$

Here $\underline{\alpha} = (\alpha_1, \dots, \alpha_L)$ are positive integers, $S_{\text{CS}}[A]$ is the Chern-Simons invariant of A , $\text{Pexp} \left(\oint_{\mathcal{L}_j} A dx \right)$ is a holonomy of A along $\mathcal{L}_j$ and Tr_{α_j} is a trace in the α_j -dimensional $SU(2)$ module. If $M = S^3$ then, according to [10],

$$J_{\underline{\alpha}}(\mathcal{L}; q) = \frac{Z_{\underline{\alpha}}(S^3, \mathcal{L}; K)}{Z(S^3; K)}, \quad Z(S^3; K) = \sqrt{\frac{2}{K}} \sin \left(\frac{\pi}{K} \right), \quad (1.80)$$

assuming that

$$q = e^{2\pi i/K}. \quad (1.81)$$

At the ‘physical level of rigor’ the path integral (1.79) can be evaluated in the stationary phase approximation as

$$K \rightarrow \infty, \quad \underline{\alpha} = \text{const.} \quad (1.82)$$

The stationary points of the Chern-Simons action are flat $SU(2)$ connections. Therefore the integral (1.79) should split into a sum of contributions of connected components $\mathcal{M}_c$ of the moduli space $\mathcal{M}$ of flat $SU(2)$ connections on M , each contribution being proportional to the ‘classical exponential’

$$Z_{\underline{\alpha}}(M, \mathcal{L}; K) = \sum_c Z_{\underline{\alpha}}^{(c)}(M, \mathcal{L}; K), \quad (1.83)$$

$$Z_{\underline{\alpha}}^{(c)}(M, \mathcal{L}; K) = e^{(iK/2\pi)S_{\text{CS}}^{(c)}} \left(\frac{4\pi^2}{K} \right)^{N_c(M)} \sum_{n=0}^{\infty} C_{\underline{\alpha};n}^{(c)}(M, \mathcal{L}) K^{-n}, \quad (1.84)$$

$$N_c(M) \in \mathbb{Q}, \quad C_{\underline{\alpha};n}^{(c)}(M, \mathcal{L}) \in \mathbb{C},$$

Here c indexes the connected components of $\mathcal{M}$ and $S_{\text{CS}}^{(c)}$ is the Chern-Simons invariant of any connection belonging to $\mathcal{M}_c$.

If a component $\mathcal{M}_c$ represents a single connection $A^{(c)}$ (that is, if $\mathcal{M}_c$ is a separate point in $\mathcal{M}$), then there are specific ‘physical’ predictions about the numbers $N_c(M)$ and $C_{\underline{\alpha};n}^{(c)}(M, \mathcal{L})$. Let D_c be a covariant derivative in the trivial $su(2)$ bundle over M associated with the connection $A^{(c)}$. $\tau_{\text{R}}^{(c)}$ denotes the Reidemeister-Ray-Singer torsion of D_c , H_c denotes the isotropy group of $A^{(c)}$ ($H_c \subset SU(2)$ commutes with the holonomies of $A^{(c)}$), $\text{Vol}(H_c)$ is the volume of H_c (if H_c is discrete, then $\text{Vol}(H_c)$ is the number of elements, otherwise the volume is calculated with the metric in which the roots of $su(2)$ have length $\sqrt{2}$) and b_M^1 is the first Betti number of M . Then, according to [10],

$$N_c(M) = \frac{1}{2} \dim H_c, \quad (1.85)$$

$$|C_{\underline{\alpha};n}^{(c)}(M, \mathcal{L})| = \frac{\prod_{j=1}^L \text{Tr}_{\alpha_j} \text{Pexp} \left(\oint_{\mathcal{L}_j} A^{(c)} dx \right)}{\text{Vol}(H_c) (\tau_{\text{R}}^{(c)})^{1/2}}. \quad (1.86)$$

If $M = S^3$, then the only flat $SU(2)$ connection on M is the trivial connection, so the sum of eq.(1.83) contains only one term

$$Z_{\underline{\alpha}}(S^3, \mathcal{L}; K) = Z_{\underline{\alpha}}^{(\text{tr})}(S^3, \mathcal{L}; K). \quad (1.87)$$

We use tr as the index c of the trivial connection. It is easy to see that

$$H_{\text{tr}} = SU(2), \quad \text{Vol}(H_{\text{tr}}) = 2^{5/2}\pi^2, \quad N_c(S^3) = 3/2, \quad \tau_{\text{R}}^{(\text{tr})} = 1, \quad (1.88)$$

so that the absolute value of the first term in the sum of eq.(1.84) is

$$\left| C_{\underline{\alpha};0}^{(\text{tr})}(S^3, \mathcal{L}) \right| = \{\underline{\alpha}\} 2^{-5/2}\pi^{-2}. \quad (1.89)$$

This is consistent with the first term of the Melvin-Morton expansion (1.46). Indeed, since

$$h = 2\pi i K^{-1} + O(K^{-3}), \quad Z(S^3; K) = 2^{1/2}\pi K^{-3/2} + O(K^{-7/2}) \quad \text{as } K \rightarrow \infty \quad (1.90)$$

then eqs.(1.80), (1.84) predict that

$$|d_{\underline{0},0}(\mathcal{L})| = \frac{K^{3/2}}{2^{1/2}\pi} \left(\frac{4\pi^2}{K} \right)^{N_c(S^3)} \left| C_{\underline{\alpha};0}^{(\text{tr})}(S^3, \mathcal{L}) \right| = 1, \quad (1.91)$$

while according to [6], $d_{\underline{0},0}(\mathcal{L}) = 1$.

The paper [3] describes a path integral representation of the colored Jones polynomial which is slightly different from eq.(1.79). Let $\mathcal{L}, \mathcal{L}'$ be a pair of links in M endowed with colors $\underline{\alpha}, \underline{\beta}$. For rational numbers $\underline{a} = (a_1, \dots, a_L)$ such that $\underline{a}/K \in \mathbb{Z}$, consider the following expression

$$Z_{\underline{\beta}}(M, \mathcal{L}, \mathcal{L}'; \underline{a}; K) = \int_{\text{b.c.}(\mathcal{L}; \underline{a})} \mathcal{D}A e^{(iK/2\pi)S_{\text{CS}}[A]} \prod_{j=1}^{L'} \text{Tr}_{\beta_j} \text{Pexp} \left(\oint_{\mathcal{L}'_j} A dx \right). \quad (1.92)$$

Here $\int_{\text{b.c.}(\mathcal{L}; \underline{a})}$ denotes a path integral over $SU(2)$ connections in the complement of the tubular neighborhood of $\mathcal{L}$ (we denote this neighborhood simply as $M \setminus \mathcal{L}$). The connections should satisfy the following boundary condition: for each connection there should exist the elements $\underline{g} \in SU(2)$, $\underline{g} = (g_1, \dots, g_L)$ such that

$$\text{Pexp} \left(\oint_{\text{mer}(\mathcal{L}_j)} A dx \right) = g_j e^{i\pi a_j \sigma_3} g_j^{-1}. \quad (1.93)$$

In eq.(1.93) we used the following notations: $\text{mer}(\mathcal{L}_j)$ is the meridian of $\mathcal{L}_j$ (that is, a simple cycle on the boundary of the tubular neighborhood of $\mathcal{L}_j$ which is contractible through that neighborhood) and σ_3 is one of the Pauli matrices:

$$\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1.94)$$

According to [3], the integral (1.92) is equal to $Z_{\underline{\alpha}, \underline{\beta}}(M, \mathcal{L} \cup \mathcal{L}'; K)$ at $\underline{a} = \underline{\alpha}/K$

$$Z_{\underline{\alpha}, \underline{\beta}}(M, \mathcal{L} \cup \mathcal{L}'; K) = Z_{\underline{\beta}}(M, \mathcal{L}, \mathcal{L}'; \underline{\alpha}/K; K). \quad (1.95)$$

Representation (1.84) coupled with eq. (1.94) allows us to apply the stationary phase approximation to the calculation of $Z_{\underline{\alpha}, \underline{\beta}}(M, \mathcal{L} \cup \mathcal{L}'; K)$ in the limit

$$K \rightarrow \infty, \quad \underline{a} = \underline{\alpha}/K = \text{const}, \quad \underline{\beta} = \text{const}. \quad (1.96)$$

The integral (1.84) should split into a sum of contributions of flat $SU(2)$ connections on $M \setminus \mathcal{L}$ which satisfy the condition (1.85).

If $M = S^3$, then among all flat connections satisfying (1.93) there are exactly 2^{L-1} $U(1)$ -reducible connections. Their holonomies belong to the same subgroup $U(1) \subset SU(2)$. These $U(1)$ -reducible connections can be indexed by a set of numbers $\underline{\mu} = (\mu_1, \dots, \mu_L)$ such that $\mu_j = \pm 1$, $2 \leq j \leq L$, $\mu_L = 2$. Using $\underline{\mu}$ as an index of the corresponding $U(1)$ -reducible connection and identifying $U(1)$ with $\{z \in \mathbb{C} \mid |z| = 1\}$ we find that

$$\text{Pexp} \left(\oint_{\text{mer}(\mathcal{L}_j)} A^{(\underline{\mu})} dx \right) = e^{i\pi \mu_j a_j}. \quad (1.97)$$

Conjecture 1.12 *The formal sum (1.50) is related to the contributions of $U(1)$ -reducible connections. Namely,*

$$\frac{Z_{\underline{\beta}}^{(\underline{\mu})}(S^3, \mathcal{L}, \mathcal{L}'; \underline{a}/K; K)}{Z(S^3; K)} = \{\underline{\mu}\} J_{\underline{\mu}, \underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q), \quad (1.98)$$

or, equivalently,

$$\frac{Z_{\underline{\beta}}^{(\underline{\mu})}(S^3, \mathcal{L}, \mathcal{L}'; \underline{a}; K)}{Z(S^3; K)} = \begin{cases} \sum_{n=0}^{\infty} P_{\underline{\beta}; n}(\mathcal{L}') h^n & \text{if } L = 0, \\ \frac{\sin(\pi a_1/K)}{\sin(\pi/K)} \sum_{n=0}^{\infty} \frac{P_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; e^{2\pi i a_1})}{\Delta_A^{2n+1}(\mathcal{L}; e^{2\pi i a_1})} h^n & \text{if } L = 1, \\ \frac{\{\underline{\mu}\}}{h} q^{\phi_0(\mathcal{L})} e^{2\pi i K \text{lk}(\mathcal{L}; \underline{\mu} \underline{a})} \sum_{n=0}^{\infty} \frac{P_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; e^{2\pi i \underline{\mu} \underline{a}})}{\nabla_A^{2n+1}(\mathcal{L}; e^{2\pi i \underline{\mu} \underline{a}})} h^n & \text{if } L \geq 2. \end{cases} \quad (1.99)$$

We have two reasons to make this conjecture. The first reason is that the first term in the sum of eq.(1.99) matches the first term of the sum of eq.(1.84) written for the $U(1)$ -reducible connection $A^{(\underline{\mu})}$. Indeed, in view of eq.(1.54), the first term in the expansion of the *r.h.s.* of eq.(1.99) for $L \geq 1$ in powers of K^{-1} becomes

$$\frac{\{\underline{\mu}\}}{2\pi i K} \frac{e^{2\pi i K \text{lk}(\mathcal{L}; \underline{\mu}\underline{a})}}{\nabla_A(\mathcal{L}; e^{2\pi i \underline{\mu}\underline{a}})} \prod_{j=1}^{L'} \frac{\sin\left(\pi \beta_j \sum_{k=1}^L \mu_k a_k\right)}{\sin\left(\pi \sum_{k=1}^L \mu_k a_k\right)}. \quad (1.100)$$

On the other hand, the Chern-Simons action of $A^{(\underline{\mu})}$ (with appropriate boundary terms included) is

$$S_{\text{CS}}^{(\underline{\mu})} = 2\pi^2 \text{lk}(\mathcal{L}; \underline{\mu}\underline{a}), \quad (1.101)$$

while the isotropy group is

$$H_{(\underline{\mu})} = U(1), \quad \text{Vol}(H_{(\underline{\mu})}) = 2^{3/2}\pi. \quad (1.102)$$

The trivial $su(2)$ bundle with connection $A^{(\underline{\mu})}$ splits into a sum of a 1-dimensional real bundle with zero connection and a 1-dimensional complex bundle with the corresponding $U(1)$ connection. The Reidemeister-Ray-Singer torsion of the former is equal to 1, while the torsion of the latter is equal to the absolute value of the Alexander-Conway function $\nabla_A(\mathcal{L}; e^{2\pi i \underline{\mu}\underline{a}})$. Therefore

$$(\tau_{\text{R}}^{(\underline{\mu})})^{1/2} = |\nabla_A(\mathcal{L}; e^{2\pi i \underline{\mu}\underline{a}})|. \quad (1.103)$$

Finally,

$$\text{Tr}_{\beta_j} \text{Pexp} \left(\oint_{\mathcal{L}'_j} A^{(\underline{\mu})} dx \right) = \frac{\sin\left(\pi \beta_j \sum_{k=1}^L \mu_k a_k\right)}{\sin\left(\pi \sum_{k=1}^L \mu_k a_k\right)}, \quad (1.104)$$

so according to eq.(1.86),

$$\left(\frac{4\pi^2}{K} \right)^{N^{(\underline{\mu})}(S^3)} \left| C_{\underline{a}, \underline{\beta}; 0}^{(\underline{\mu})}(S^3, \mathcal{L} \cup \mathcal{L}') \right| = \frac{(2K)^{-1/2}}{|\nabla_A(\mathcal{L}; e^{2\pi i \underline{\mu}\underline{a}})|} \prod_{j=1}^{L'} \left| \frac{\sin\left(\pi \beta_j \sum_{k=1}^L \mu_k a_k\right)}{\sin\left(\pi \sum_{k=1}^L \mu_k a_k\right)} \right|. \quad (1.105)$$

If we substitute eqs. (1.101) and (1.105) into the expression (1.84), then we can see easily that the dominant terms of the K^{-1} expansion of both sides of eq. (1.98) coincide at least up to a K -independent phase factor which may come from $C_{\underline{\alpha}, \underline{\beta}; 0}^{(\mu)}(S^3, \mathcal{L} \cup \mathcal{L}')$.

The second reason for making Conjecture 1.12 is the relation between $J_{\underline{\alpha}, \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q)$ and $J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q)$ established in Corollary 1.8. Its path integral interpretation relies on the following theorem.

For positive numbers $\underline{c} = (c_1, \dots, c_L)$ and positive integers $\underline{n} = (n_1, \dots, n_{L-1})$ let $\mathcal{A}_{\underline{c}, \underline{n}}$ denote a set of all points $\underline{a} = (a_1, \dots, a_L)$ which satisfy the inequalities

$$a_1 < c_1, \quad a_{j+1} < c_{j+1} a_j^{n_j}, \quad 1 \leq j \leq L-1, \quad a_L > 0. \quad (1.106)$$

Theorem 1.13 *Let $\mathcal{L} \subset S^3$ be an algebraically connected link with admissible indexing. Then there exist positive numbers $\underline{x} = (x_1, \dots, x_L)$ and positive integers $\underline{n} = (n_1, \dots, n_{L-1})$ such that for any $\underline{a} \in \mathcal{A}_{\underline{c}, \underline{n}}$ the only flat connections in the link complement which satisfy conditions (1.93) are the $U(1)$ -reducible ones.*

We will prove this theorem in Appendix A.

Theorem 1.13 implies that $U(1)$ -reducible connections $A^{(\mu)}$ are the only contributors to $Z_{\underline{\beta}}(S^3, \mathcal{L}, \mathcal{L}'; \underline{a}; K)$ within the domain (1.106). In other words, in that domain

$$Z_{\underline{\beta}}(S^3, \mathcal{L}, \mathcal{L}'; \underline{a}; K) = \sum_{\substack{\mu_j = \pm 1 \\ 2 \leq j \leq L \\ \mu_1 = 1}} Z_{\underline{\beta}}^{(\mu)}(S^3, \mathcal{L}, \mathcal{L}'; \underline{a}; K) \quad (1.107)$$

in the limit (1.96). Now we want to pass from the limit (1.96) to the limit

$$K \rightarrow \infty, \quad \underline{\alpha}, \underline{\beta} = \text{const.} \quad (1.108)$$

This limit is similar to eq. (1.82), so $Z_{\underline{\alpha}, \underline{\beta}}(S^3, \mathcal{L} \cup \mathcal{L}'; K)$ receives the contribution only from the trivial connection on S^3 , and this contribution is of the form

$$\begin{aligned} Z_{\underline{\alpha}, \underline{\beta}}(S^3, \mathcal{L} \cup \mathcal{L}'; K) &= Z_{\underline{\alpha}, \underline{\beta}}^{(\text{tr})}(S^3, \mathcal{L} \cup \mathcal{L}'; K) \\ &= Z(S^3; K) \sum_{\substack{\underline{m}, \underline{m}', n \geq 0 \\ |\underline{m}| + |\underline{m}'| \leq n}} d_{\underline{m}, \underline{m}'; n}(\mathcal{L} \cup \mathcal{L}') \underline{\alpha}^{2\underline{m}+1} \underline{\beta}^{2\underline{m}'+1} h^n \end{aligned} \quad (1.109)$$

(recall that in view of the first of eqs. (1.90) expansion in powers of h is equivalent to expansion in powers of K^{-1}). A natural way to pass from eq. (1.107) to eq. (1.109) is to expand $Z_{\underline{\beta}}^{(\underline{\mu})}(S^3, \mathcal{L}, \mathcal{L}'; \underline{a}; K)$ in powers of $\underline{a}$ while substituting $\underline{a} = \underline{\alpha}/K$. However, we have to stay within the domain (1.106) while performing this expansion. This can be done by expanding in powers of $\underline{a}$ in sequence: starting with a_L and ending with a_1 . Therefore we conclude that

$$Z_{\underline{\alpha}, \underline{\beta}}(S^3, \mathcal{L} \cup \mathcal{L}'; K) = \left[\sum_{\substack{\mu_j = \pm 1 \\ 2 \leq j \leq L \\ \mu_1 = 1}} Z_{\underline{\beta}}^{(\underline{\mu})}(S^3, \mathcal{L}, \mathcal{L}'; \underline{a}/K; K) \right]_{(\underline{\alpha})}. \quad (1.110)$$

Thus the second reason for making Conjecture 1.12 is that relation (1.98) makes eq. (1.110) equivalent to eq. (1.68).

2 Calculating the RC invariant

The best way to derive $J_{\underline{\alpha}, \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q)$ would be to study the asymptotic behaviour of the colored Jones polynomial $J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q)$ in the limit (1.96) or, in other words, in the limit

$$q \rightarrow 1, \quad q^{\underline{\alpha}}, \underline{\beta} = \text{const.} \quad (2.1)$$

Since we can not do this directly yet, we have to take a detour. We will study the limit

$$q \rightarrow 1, \quad \underline{\alpha}, \underline{\beta} = \text{const.} \quad (2.2)$$

This means that we will work with the Melvin-Morton expansion

$$J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q) = \sum_{n=0}^{\infty} P_{\underline{\alpha}, \underline{\beta}; n}(\mathcal{L} \cup \mathcal{L}') h^n, \quad P_{\underline{\alpha}, \underline{\beta}; n}(\mathcal{L} \cup \mathcal{L}') \in \mathbb{Q}[\underline{\alpha}, \underline{\beta}] \quad (2.3)$$

as it comes from the $\check{R}$ -matrix representation described in subsection 1.2.7. However we will derive the expansion (2.3) in two stages. At the first stage we will expand all instances of q in $\check{R}$ and Q_q except for the expressions of the form

$$q^{\alpha_i \alpha_j}, \quad q^{\alpha_j}, \quad 1 \leq i, j \leq L, \quad (2.4)$$

which we will leave intact. This way we will present $J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q)$ in the form which is very close to eq.(1.65). At the second stage we will expand the expressions (2.4) thus coming to the Melvin-Morton expansion (2.3) in the form (1.68).

2.1 Preliminary calculations

2.1.1 Breaking a strand

We start with eq.(1.44), according to which

$$J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q) = q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')} \text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}} Q_q B_{\mathcal{L} \cup \mathcal{L}'}, \quad (2.5)$$

where $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ is an N -strand braid whose closure produces the link $\mathcal{L} \cup \mathcal{L}'$.

Let us introduce a notation

$$\alpha_* = \begin{cases} \alpha_1 & \text{if } \mathcal{L} \text{ is non-empty} \\ \beta_1 & \text{if } \mathcal{L} \text{ is empty} \end{cases} \quad (2.6)$$

The space $V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}$ is a tensor product

$$V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}} = V_{\alpha_*} \otimes V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}^{[1]}, \quad (2.7)$$

where $V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}^{[1]}$ is what remains of the tensor product $V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}$ after we remove the first factor V_{α_*} . We introduce a modified version of the operator (1.43)

$$Q_q^{[1]} = I \otimes (q^{H/2})^{\otimes (N-1)}. \quad (2.8)$$

Obviously,

$$\text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}} Q_q B_{\mathcal{L} \cup \mathcal{L}'} = \text{Tr}_{V_{\alpha_*}} q^{H/2} \text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}^{[1]}} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'}. \quad (2.9)$$

On the other hand, since the operator $B_{\mathcal{L} \cup \mathcal{L}'}$ commutes with the generators of $SU_q(2)$ and since the module V_{α_*} is irreducible, then the operator $\text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}^{[1]}} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'}$ is proportional to I

$$\text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}^{[1]}} Q_q B_{\mathcal{L} \cup \mathcal{L}'} = CI, \quad C \in \mathbb{C}. \quad (2.10)$$

Therefore since $\text{Tr}_{V_{\alpha_*}} q^{H/2} = [\alpha_*]$, then

$$\text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}} Q_q B_{\mathcal{L} \cup \mathcal{L}'} = [\alpha_*] C. \quad (2.11)$$

There are various ways to determine the constant C . For example, since

$$\lambda_0^{\alpha_*-1} \text{Tr}_{V_{\alpha_*}} \lambda_0^{-H} \Big|_{\lambda_0=0} = 1, \quad (2.12)$$

then

$$C = \lambda_0^{\alpha_*-1} \text{Tr}_{V_{\alpha_*}} \lambda_0^{-H} \text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}}^{[1]} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'} \Big|_{\lambda_0=0}, \quad (2.13)$$

and combining eqs. (1.44) and (2.11) we conclude that

$$J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q) = q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')} [\alpha_*] \lambda_0^{\alpha_*-1} \text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}} (\lambda_0^{-H} \otimes I^{\otimes(N-1)}) Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'} \Big|_{\lambda_0=0}. \quad (2.14)$$

2.1.2 Twisting $\check{R}$ -matrix

In subsection 1.2.7 we described a representation of the braid group in which a positive elementary braid is mapped into the matrix $\check{R} = PR$. It is better for our purposes to use a slightly modified representation in which the image of a positive elementary braid is defined as

$$\check{R} = q^{-\frac{3}{8}(\hat{\gamma}-1) \otimes H - \frac{1}{8}H \otimes (\hat{\gamma}-1)} PR q^{\frac{3}{8}(\hat{\gamma}-1) \otimes H + \frac{1}{8}H \otimes (\hat{\gamma}-1)}. \quad (2.15)$$

Here $\hat{\gamma}$ is an operator which multiplies the elements of the module V_γ by γ

$$\hat{\gamma}: V_\gamma \rightarrow V_\gamma, \quad \hat{\gamma}(f_m) = \gamma f_m. \quad (2.16)$$

As a result, the matrices $\check{R}$ and $\check{R}^{-1}$ map $V_{\gamma_1} \otimes V_{\gamma_2}$ into $V_{\gamma_2} \otimes V_{\gamma_1}$ according to the formulas

$$\check{R}(f_{m_1} \otimes f_{m_2}) = \sum_{0 \leq n \leq \max\{m_1, \gamma_2 - m_2 - 1\}} \check{R}_{m_1, m_2}^{m_2+n, m_1-n} f_{m_2+n} \otimes f_{m_1-n}, \quad (2.17)$$

$$\check{R}^{-1}(f_{m_1} \otimes f_{m_2}) = \sum_{0 \leq n \leq \max\{m_2, \gamma_1 - m_1 - 1\}} (\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} f_{m_2-n} \otimes f_{m_1+n}, \quad (2.18)$$

where

$$\check{R}_{m_1, m_2}^{m_2+n, m_1-n} = q^{(\gamma_1-1)(\gamma_2-1)/4} \prod_{l=1}^n \left(\frac{(q^{-m_2-l} - q^{-\gamma_2})(q^{m_1-n+l} - 1)}{q^l - 1} \right) \times q^{-m_2(\gamma_1-1)+m_1m_2+\frac{1}{2}n(n+1)}, \quad (2.19)$$

$$(\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} = q^{-(\gamma_1-1)(\gamma_2-1)/4} \prod_{l=1}^n \left(\frac{(q^{m_1+l} - q^{\gamma_1})(q^{-m_2+n-l} - 1)}{q^{-l} - 1} \right) \times q^{m_1(\gamma_2-1)+n(\gamma_2-\gamma_1)-m_1m_2-\frac{1}{2}n(n+1)}. \quad (2.20)$$

The reason for twisting $\check{R}$ -matrix is that the matrix elements (2.19) and (2.20) have certain integrality properties.

Lemma 2.1 *There exist the polynomials $S^{(\pm)}(t_1, t_2, q) \in \mathbb{Z}[t_1, t_1^{-1}, t_2, t_2^{-1}, q, q^{-1}]$ such that*

$$\check{R}_{m_1, m_2}^{m_2+n, m_1-n} = q^{(\gamma_1-1)(\gamma_2-1)/4} S^{(+)}(q^{\alpha_1}, q^{\alpha_2}, q), \quad (2.21)$$

$$(\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} = q^{-(\gamma_1-1)(\gamma_2-1)/4} S^{(-)}(q^{\alpha_1}, q^{\alpha_2}, q). \quad (2.22)$$

Proof. Since

$$\prod_{l=1}^n \left(\frac{q^{m_1-n+l} - 1}{q^l - 1} \right), \quad \prod_{l=1}^n \left(\frac{q^{-m_2+n-l} - 1}{q^{-l} - 1} \right), \quad q^{\pm \frac{1}{2}n(n+1)} \in \mathbb{Z}[q, q^{-1}], \quad (2.23)$$

then we can simply choose

$$S^{(+)}(t_1, t_2, q) = \prod_{l=1}^n \left(\frac{(q^{-m_2-l} - t_2^{-1})(q^{m_1-n+l} - 1)}{q^l - 1} \right) t_1^{-m_2} q^{m_2+m_1m_2+\frac{1}{2}n(n+1)}, \quad (2.24)$$

$$S^{(-)}(t_1, t_2, q) = \prod_{l=1}^n \left(\frac{(q^{m_1+l} - t_1)(q^{-m_2+n-l} - 1)}{q^{-l} - 1} \right) t_1^{-n} t_2^{m_1+n} q^{-m_1-m_1m_2-\frac{1}{2}n(n+1)}. \quad (2.25)$$

□

From now on we will use the matrices (2.19), (2.20) instead of PR and its inverse in order to construct the operator $B_{\mathcal{L} \cup \mathcal{L}'}$ which represents the braid $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$. However, this substitution does not affect eq.(2.14). Indeed, if $V_{\gamma_1} \otimes V_{\gamma_2}$ is a part of the tensor product $V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}$, then a conjugation of the matrix PR by the operator $q^{\frac{3}{8}(\hat{\gamma}-1) \otimes H + \frac{1}{8}H \otimes (\hat{\gamma}-1)}$ in eq.(2.15) is equivalent to the conjugation of the whole space $V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}$ by

$$\bigotimes_{i=1}^N q^{\frac{H}{8} (3 \sum_{j=1}^{i-1} (\gamma(j)-1) + \sum_{j=i+1}^N (\gamma(j)-1))}, \quad (2.26)$$

where γ maps the number of the strand to the color of the corresponding link component. Thus we see that the conjugation in eq.(2.15) has no effect on the trace of eq.(2.14).

2.1.3 Parametrized and regularized trace

We want to introduce two more operators acting on $V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}$. For a positive integer N and for two complex variables $\underline{\lambda} = (\lambda_0, \lambda)$ we introduce an operator

$$Q_{\underline{\lambda}}^{(N)} = \lambda_0^{-H} \otimes (\lambda^{-H})^{\otimes (N-1)} \quad (2.27)$$

and a parametrized trace

$$J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q | \underline{\lambda}) = q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')} [\alpha_*] \text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}} Q_{\underline{\lambda}}^{(N)} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'}. \quad (2.28)$$

In contrast to the Jones polynomial, the parametrized trace (2.28) is not an invariant of the link $\mathcal{L} \cup \mathcal{L}'$, because it depends on the choice of the braid $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ whose closure is $\mathcal{L} \cup \mathcal{L}'$. Equation (2.14) implies a relation between the parametrized trace and the colored Jones polynomial

$$\lambda_0^{\alpha_* - 1} J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q | \underline{\lambda}) \Big|_{\substack{\lambda=1 \\ \lambda_0=0}} = J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q). \quad (2.29)$$

Next we introduce a projection operator acting on V_γ

$$P_M(f_m) = \begin{cases} f_m & \text{if } |\gamma - 1 - 2m| \leq M, \\ 0 & \text{if } |\gamma - 1 - 2m| > M. \end{cases} \quad (2.30)$$

Let us replace the operators $\check{R}$ in the expression for $B_{\mathcal{L} \cup \mathcal{L}'}$ with ‘regularized’ operators $(P_M \otimes P_M) \check{R} (P_M \otimes P_M)$ at $(\mathcal{L}', \mathcal{L}')$ crossings, with operators $(P_M \otimes I) \check{R} (I \otimes P_M)$ at $(\mathcal{L}, \mathcal{L}')$ crossings and with operators $(I \otimes P_M) \check{R} (P_M \otimes I)$ at $(\mathcal{L}', \mathcal{L})$ crossings. We perform the same substitution for the operators $\check{R}^{-1}$. We denote the resulting regularized braid operator as $B_{\mathcal{L} \cup \mathcal{L}'; M}$. We also introduce a projection operator

$$P_{\mathcal{L} \cup \mathcal{L}'; M} = P_M^{\otimes N} \quad (2.31)$$

and we define a parametrized regularized trace as

$$J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q | \underline{\lambda}; M) = q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')} [\alpha_*] \text{Tr}_{V_{\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'; \underline{\alpha}, \underline{\beta}}}} P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda}}^{(N)} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M}. \quad (2.32)$$

Since $P_M = I$ on V_γ if $M \geq \gamma - 1$, then

$$J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q | \underline{\lambda}; M) = J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q | \underline{\lambda}) \quad \text{if } \underline{M} \geq \max\{\underline{\alpha}, \underline{\beta}\} - 1. \quad (2.33)$$

2.1.4 Verma module resolution

Our next step is to resolve the traces over $SU_q(2)$ modules attached to the strands of $\mathcal{L}$ in terms of Verma modules. For $\gamma \in \mathbb{Z}$, an $SU_q(2)$ Verma module $V_{\gamma, \infty}$ is an infinite-dimensional space with the basis f_m , $m \geq 0$. The action of the $SU_q(2)$ generators on $V_{\gamma, \infty}$ is given by eqs.(1.34)–(1.36), while the operators $\check{R}$ and $\check{R}^{-1}$ map the tensor product $V_{\gamma_1, \infty} \otimes V_{\gamma_2, \infty}$ into $V_{\gamma_2, \infty} \otimes V_{\gamma_1, \infty}$ according to the formulas which are similar to (2.17), (2.18)

$$\check{R}(f_{m_1} \otimes f_{m_2}) = \sum_{0 \leq n \leq m_1} \check{R}_{m_1, m_2}^{m_2+n, m_1-n} f_{m_2+n} \otimes f_{m_1-n}, \quad (2.34)$$

$$\check{R}^{-1}(f_{m_1} \otimes f_{m_2}) = \sum_{0 \leq n \leq m_2} (\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} f_{m_2-n} \otimes f_{m_1+n}, \quad (2.35)$$

where the matrix elements are given by eqs.(2.19) and (2.20).

For a positive integer γ there is a natural immersion

$$V_\gamma \rightarrow V_{\gamma, \infty}, \quad f_m \mapsto f_m. \quad (2.36)$$

Moreover, there is an isomorphism of $SU_q(2)$ modules

$$V_{\gamma, \infty}/V_\gamma = V_{-\gamma, \infty}. \quad (2.37)$$

If we choose the vectors f_m , $m \geq \gamma$ as the basis of $V_{\gamma, \infty}/V_\gamma$, then this isomorphism identifies

$$f_{m+n} \leftrightarrow \frac{[\gamma + m]!}{[m]!} f_m. \quad (2.38)$$

We need another family of infinite-dimensional modules. For an integer γ , $V_{\gamma, \pm\infty}$ is an infinite-dimensional space with the basis f_m , $m \in \mathbb{Z}$. The action of the $SU_q(2)$ generators

on $V_{\gamma,\infty}$ is given again by eqs. (1.34)–(1.36), while the operators $\check{R}$ and $\check{R}^{-1}$ map the tensor product $V_{\gamma_1,\pm\infty} \otimes V_{\gamma_2,\pm\infty}$ into $V_{\gamma_2,\pm\infty} \otimes V_{\gamma_1,\pm\infty}$ according to the formulas which are similar to (2.17), (2.18)

$$\check{R}(f_{m_1} \otimes f_{m_2}) = \sum_{n \geq 0} \check{R}_{m_1, m_2}^{m_2+n, m_1-n} f_{m_2+n} \otimes f_{m_1-n}, \quad (2.39)$$

$$\check{R}^{-1}(f_{m_1} \otimes f_{m_2}) = \sum_{n \geq 0} (\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} f_{m_2-n} \otimes f_{m_1+n}, \quad (2.40)$$

where the matrix elements are given again by eqs. (2.19) and (2.20). Also, for a positive γ there is an immersion

$$V_\gamma \rightarrow V_{\gamma,\pm\infty}, \quad f_m \mapsto f_m. \quad (2.41)$$

Consider two tensor products of $SU_q(2)$ modules

$$V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha},\infty} = \bigotimes_{j=1}^L V_{\alpha_j,\infty}^{\otimes N_j}, \quad V_{\mathcal{B}_{\mathcal{L}'};\underline{\beta},\pm\infty} = \bigotimes_{j=1}^{L'} V_{\beta_j,\pm\infty}, \quad (2.42)$$

where in the second tensor product

$$V_{\mathcal{B}_{\mathcal{L}'};\beta_j,\pm\infty} = V_{\beta_j} \otimes V_{\beta_j,\pm\infty}^{\otimes (N'_j-1)}. \quad (2.43)$$

Since the action of $\check{R}$ -matrix was extended to spaces $V_{\gamma,\infty}$ and $V_{\gamma,\pm\infty}$, then the action of the operator $Q_{\underline{\lambda}} Q_q^{[1]} B_M$ can be extended to $V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha},\infty} \otimes V_{\mathcal{B}_{\mathcal{L}'};\underline{\beta}}$ and $V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha},\infty} \otimes V_{\mathcal{B}_{\mathcal{L}'};\underline{\beta},\pm\infty}$. For a set of integer (not necessarily positive) colors $\underline{\alpha}$ and for a set of positive integer colors $\underline{\beta}$ we define the following trace

$$J_{\underline{\alpha};\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')} [\alpha_*] \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha},\infty} \otimes V_{\mathcal{B}_{\mathcal{L}'};\underline{\beta}}} P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda}}^{(N)} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M}. \quad (2.44)$$

Proposition 2.2 *The trace $J_{\underline{\alpha};\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$ has the following properties:*

(1) *If $\mathcal{L}$ is empty, then*

$$J_{\underline{\beta}}^{(r)}(\emptyset, \mathcal{L}'; q | \underline{\lambda}; M) = J_{\underline{\beta}}(\mathcal{L}'; q | \underline{\lambda}; M). \quad (2.45)$$

(2) For any j , $1 \leq j \leq L'$,

$$J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) \Big|_{\beta_j=1} = J_{\underline{\alpha}; \underline{\beta}_{(j)}}^{(r)}(\mathcal{L}, \mathcal{L}'_{(j)}; q | \underline{\lambda}; M). \quad (2.46)$$

(3) The space $V_{\mathcal{B}_{L'}; \underline{\beta}}$ can be replaced by $V_{\mathcal{B}_{L'}; \underline{\beta}, \pm\infty}$ in eq. (2.44)

$$J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')} [\alpha_*] \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{L'}; \underline{\beta}, \pm\infty}} P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda}}^{(N)} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M} \quad (2.47)$$

(4) If $\mathcal{L}$ is non-empty, then for any j , $1 \leq j \leq L$

$$J_{\underline{\alpha}_{(j)}; \underline{\beta}_0, \underline{\beta}}^{(r)}(\mathcal{L}_{(j)}, \mathcal{L}_j \cup \mathcal{L}'; q | \underline{\lambda}; M) = \sum_{\mu=\pm 1} \mu J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) \Big|_{\alpha_j=\mu\beta_0}. \quad (2.48)$$

Proof of claims 1 and 2 of Proposition 2.2. Equation (2.46) is obvious if we compare eqs. (2.44) and (2.32). To prove eq. (2.57) we observe that $\dim V_{\beta_{L'}} = 1$ if $\beta_{L'} = 1$. As a result, the matrix elements of $\check{R}$ -matrices (2.17), (2.18) which involve at least one space $V_{\beta_{L'}}$ are equal to 1. The same is true for the matrix elements of the operators $P_{\mathcal{L} \cup \mathcal{L}'; M}$, $Q_{\underline{\lambda}}^{(N)}$ and $Q_q^{[1]}$ which act on $V_{\beta_{L'}}$. Finally, it follows from eq. (1.45) that

$$q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')} \Big|_{\beta_{L'}=1} = q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}'_{[1]})} \quad (2.49)$$

and thus we come to eq. (2.46). $\square$

The proof of claims 3 and 4 of Proposition 2.2 requires a simple lemma

Lemma 2.3 *Let V be a finite-dimensional linear space with a subspace $W \subset V$. For a fixed positive integer N we denote*

$$U_j = V^{\otimes(j-1)} \otimes W \otimes V^{\otimes(N-j)}, \quad 1 \leq j \leq N. \quad (2.50)$$

Suppose that a linear operator O acting on the space $V^{\otimes N}$ has the following property: there exists a cyclic permutation of N elements σ such that

$$O(U_j) \subset U_{\sigma(j)} \quad \text{for any } j, 1 \leq j \leq N. \quad (2.51)$$

Then

(1) The action of O is well-defined on the spaces $W^{\otimes N}$, $(V/W)^{\otimes N}$ and there is a simple trace formula

$$\mathrm{Tr}_{W^{\otimes N}} O = \mathrm{Tr}_{V^{\otimes N}} O - \mathrm{Tr}_{(V/W)^{\otimes N}} O. \quad (2.52)$$

(2) Let j be a fixed number, $1 \leq j \leq N$. A partial trace of O over $N - 1$ spaces V of the tensor product $V^{\otimes N}$ except the first space V defines an operator $\mathrm{Tr}_{V^{\otimes(N-1)}} O$ acting on V . Then $\mathrm{Tr}_{V^{\otimes(N-1)}}(W) \subset W$ and

$$\mathrm{Tr}_{W^{\otimes N}} O = \mathrm{Tr}_{U_1} O. \quad (2.53)$$

Proof. We leave the proof to the reader. $\square$

Proof of claims 3 and 4 of Proposition 2.2. Since the R -matrix, as well as the operators P_M , $Q_{\underline{\lambda}}^{(N)}$ and $Q_q^{[1]}$ can be expressed in terms of the generators of $SU_q(2)$, then their action commutes with immersions (2.36) and (2.41). Upon being acted on by the $SU_q(2)$ generators of R -matrices, the elements of $SU_q(2)$ modules are directed along the braid strands by the permutation operators P which are present at every elementary crossing. The permutation generated by B among the strands of any particular link component is cyclic. Therefore it is easy to see that the operator $Q_{\underline{\lambda}}^{(N)} Q_q^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M}$ satisfies the conditions of Lemma 2.3 for any space $V_{\alpha_j, \infty}^{\otimes N_j}$ (V being $V_{\alpha_j, \infty}$ and W being V_{α_j}) and for any space $V_{\beta_j, \pm\infty}^{\otimes N'_j}$ (V being $V_{\beta_j, \pm\infty}$ and W being V_{β_j}). The infinite dimension of the spaces $V_{\alpha_j, \infty}$ and $V_{\beta_j, \pm\infty}$ is not important, since the presence of the projection operators P_M in the expression for $B_{\mathcal{L} \cup \mathcal{L}'; M}$ makes the calculation of the traces effectively finite-dimensional. Therefore eq.(2.47) follows from eq.(2.53), and eq.(2.48) follows from eq.(2.52). $\square$

Proposition 2.2 has the following

Corollary 2.4 *The regularized trace (2.32) can be presented as a linear combination of the traces (2.44)*

$$J_{\underline{\alpha}, \underline{\beta}}(\mathcal{L} \cup \mathcal{L}'; q | \underline{\lambda}; M) = \sum_{\substack{\mu_j = \pm 1 \\ 1 \leq j \leq L}} \{ \underline{\mu} \} J_{\underline{\mu}, \underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L} \cup \mathcal{L}'; q | \underline{\lambda}; M), \quad (2.54)$$

Proof. Equation (2.54) is a result of applying eq.(2.48) consequently to all the components of the link $\mathcal{L}$. $\square$

2.2 Expansion of parametrized trace in powers of h

Our next step is to perform a partial expansion of the trace $J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$ in powers of h . We want to expand all instances of q except for the expressions (2.4).

2.2.1 General properties

Let us first establish some properties of the trace $J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$. We introduce the modified operators

$$Q_{q,*}^{[1]} = I \otimes (q^{-\hat{m}})^{\otimes(N-1)}, \quad Q_{\underline{\lambda},*}^{(N)} = \lambda_0^{2\hat{m}} \otimes (\lambda^{2\hat{m}})^{\otimes(N-1)}. \quad (2.55)$$

We also introduce a modified operator $B_{\mathcal{L} \cup \mathcal{L}'; M, *}$ which is constructed in the same way as $B_{\mathcal{L} \cup \mathcal{L}'; M}$, except that we drop the factors $q^{\pm(\gamma_1-1)(\gamma_2-1)/4}$ from the matrix elements of $\check{R}$ and $\check{R}^{-1}$.

Proposition 2.5 *For a given braid $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ and for fixed M and $\underline{\beta}$ there exists a unique polynomial*

$$A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q | M) \in \mathbb{Z}[\underline{\lambda}, \underline{t}, \underline{t}^{-1}, q, q^{-1}]$$

such that

$$J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = [\alpha_*] q^{\phi(\mathcal{L} \cup \mathcal{L}' | \underline{\alpha}, \underline{\beta})} \Lambda(\underline{\alpha}, \underline{\beta}; \underline{\lambda}) A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; q^{\underline{\alpha}}; q | M), \quad (2.56)$$

$$A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; q^{\underline{\alpha}}; q | M) = \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}} P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda},*}^{(N)} Q_{q,*}^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M, *}, \quad (2.57)$$

where

$$\begin{aligned} \phi(\mathcal{L} \cup \mathcal{L}' | \underline{\alpha}, \underline{\beta}) &= -\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}') \\ &+ \frac{1}{4} \left(\sum_{1 \leq i, j \leq L} l_{ij}(\alpha_i - 1)(\alpha_j - 1) + \sum_{1 \leq i, j \leq L'} l'_{ij}(\beta_i - 1)(\beta_j - 1) + 2 \sum_{\substack{1 \leq i \leq L \\ 1 \leq j \leq L'}} l''_{ij}(\alpha_i - 1)(\beta_j - 1) \right) \\ &+ \frac{1}{2} \left(\sum_{j=1}^L N_j(\alpha_j - 1) + \sum_{j=1}^{L'} N'_j(\beta_j - 1) + (1 - \alpha_*) \right) \end{aligned} \quad (2.58)$$

and

$$\Lambda(\underline{\alpha}, \underline{\beta}; \underline{\lambda}) = \lambda_0^{1-\alpha_*} \lambda^{\sum_{j=1}^L N_j(1-\alpha_j) + \sum_{j=1}^{L'} N'_j(1-\beta_j) + (\alpha_*-1)}. \quad (2.59)$$

Proof. Let us review the structure of the matrix elements of the operators $Q_{\underline{\lambda}}^{(N)}$, $Q_q^{[1]}$ and $B_{\mathcal{L} \cup \mathcal{L}'; M}$ in the basis f_m . For $f_m \in V_\gamma$

$$\lambda^{-H}(f_m) = \lambda^{1-\gamma} \lambda^{2m} f_m, \quad q^{H/2}(f_m) = q^{(\gamma-1)/2} q^{-m} f_m. \quad (2.60)$$

The expression for $B_{\mathcal{L} \cup \mathcal{L}'; M}$ contains the matrices $\check{R}$ and $\check{R}^{-1}$, whose matrix elements are described by Lemma 2.1. Thus, if we assemble the factors $\lambda^{1-\gamma}$ of the first of eqs.(2.60) into $\Lambda(\underline{\alpha}, \underline{\beta}; \underline{\lambda})$ and if we assemble the factors $q^{(\gamma-1)/2}$ of the second of eqs.(2.60) together with the factors $q^{\pm(\gamma_1-1)(\gamma_2-1)/4}$ of eqs.(2.19), (2.20) and with the factor $q^{-\phi_{\text{sl}}(\mathcal{L} \cup \mathcal{L}')}$ of eq.(2.47) into the factor $q^{\phi(\mathcal{L} \cup \mathcal{L}' | \underline{\alpha}, \underline{\beta})}$, then the rest of the product of matrix elements of the operators $\check{R}$, $\check{R}^{-1}$, P_M , $Q_{\underline{\lambda}}^{(N)}$ and $Q_q^{[1]}$ is a polynomial of q^α , $q^{-\alpha}$, and q, q^{-1} with integer coefficients. Note that if we use either eq.(2.54) or eq.(2.47) for $J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$, then due to the presence of projection operators P_M , the range of summation of the indices of the basis vectors $f_{m_1} \otimes \cdots \otimes f_{m_N}$ used to calculate the matrix elements is finite, and the boundaries of this range do not depend on $\underline{\alpha}$. Thus, taking a sum of the product of matrix elements over the indices of basis vectors we come to eq.(2.56). The uniqueness of the polynomial $A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q | M)$ which satisfies eq.(2.47) for any $\underline{\alpha}$ is obvious. $\square$

If we expand the polynomials $A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q | M)$ in powers of $h = q - 1$ then we come to the following

Proposition 2.6 *For a braid $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ whose closure is $\mathcal{L} \cup \mathcal{L}'$, there is a unique set of polynomials*

$$A_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t} | M) \in \mathbb{Z}[\underline{\lambda}, \underline{t}, \underline{t}^{-1}] \quad (2.61)$$

such that for fixed $\underline{\alpha}, \underline{\beta}$ and M the following series in powers of h converges in the vicinity of $h = 0$

$$J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = [\alpha_*] q^{\phi(\mathcal{L} \cup \mathcal{L}' | \underline{\alpha}, \underline{\beta})} \Lambda(\underline{\alpha}, \underline{\beta}; \underline{\lambda}) \sum_{n=0}^{\infty} A_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; q^\alpha | M) h^n. \quad (2.62)$$

The polynomials $A_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t} | M)$ satisfy the following property: for any $k \geq 0$

$$\Phi_2(\alpha_i; \underline{t}_{(i)}; \underline{\lambda}; \underline{\beta}) \sum_{n=0}^k A_{\alpha_i, \underline{\beta}; n}(\mathcal{L}_{(i)}, \mathcal{L}_i \cup \mathcal{L}'; \underline{\lambda}; \underline{t}_{(i)} | M) \quad (2.63)$$

$$= \sum_{\mu=\pm 1} \mu \Phi_2(\mu\alpha_i; \underline{t}_{(i)}; \underline{\lambda}; \underline{\beta}) \sum_{n=0}^k A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M) \Big|_{t_i=q^{\mu\alpha_i}} + O(h^{k+1}) \quad \text{as } h \rightarrow 0,$$

where

$$\Phi_2(\alpha_i; \underline{t}_{(i)}; \underline{\lambda}; \underline{\beta}) = \lambda_0^{-\delta_{1i}\alpha_i} \lambda^{(\delta_{1i}-N_i)\alpha_i} q^{\frac{\alpha_i}{2}(\sum_{j=1}^{L'} l''_{ij}(\beta_i-1) - \sum_{j=1}^L l_{ij} + N_i - \delta_{1i})} \prod_{\substack{1 \leq j \leq L \\ j \neq i}} t_j^{l_{ij}\alpha_i/2} \quad (2.64)$$

and δ_{1i} is the Kronecker symbol.

Proof. The uniqueness of $A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M)$ is obvious. A set of polynomials satisfying eq.(2.62) can be obtained by expanding the polynomials $A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q|M)$ in powers of h

$$A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q|M) = \sum_{n=0}^{\infty} A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M) h^n, \quad A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M) \in \mathbb{Z}[\underline{\lambda}, \underline{t}, \underline{t}^{-1}], \quad (2.65)$$

Now we prove eq.(2.63). It follows from eqs.(2.48), (2.56) and from the uniqueness of the polynomials $A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q|M)$ that

$$\begin{aligned} \Phi_2(\alpha_i; \underline{t}_{(i)}; \underline{\lambda}; \underline{\beta}) A_{\alpha_i, \underline{\beta}}(\mathcal{L}_{(i)}, \mathcal{L}_i \cup \mathcal{L}'; \underline{\lambda}; \underline{t}_{(i)}; q|M) \\ = \sum_{\mu=\pm 1} \mu \Phi_2(\mu\alpha_i; \underline{t}_{(i)}; \underline{\lambda}; \underline{\beta}) A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q|M) \Big|_{t_i=q^{\mu\alpha_i}}. \end{aligned} \quad (2.66)$$

If we expand the polynomials A in both sides of this equation in powers of h , then we come to eq.(2.63). $\square$

It turns out that the polynomials $A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M)$ have a limit as $M \rightarrow \infty$.

Proposition 2.7 *For a braid $\mathcal{B}$ whose closure is $\mathcal{L} \cup \mathcal{L}'$ there exist the polynomials*

$$Q(\mathcal{L}; \underline{t}; \underline{\lambda}), Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}) \in \mathbb{Z}[\underline{t}, \underline{t}^{-1}, \underline{\lambda}] \quad (2.67)$$

such that there is a vicinity of $\underline{t} = 1$, $\underline{\lambda} = 0$ in which for fixed $\underline{\beta}$ there exists a limit

$$\lim_{M \rightarrow \infty} A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M) = \frac{Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda})}{Q^{2n+1}(\mathcal{L}; \underline{t}; \underline{\lambda})}. \quad (2.68)$$

The polynomials (2.67) can be selected in such a way that

$$Q(\mathcal{L}; \underline{t}; \underline{\lambda}) \Big|_{\substack{\lambda_0=0 \\ \lambda=1}} = \begin{cases} \Phi_1^{-1}(\mathcal{B}_{\mathcal{L}}; \underline{t}) (t_1^{1/2} - t_1^{-1/2}) \nabla_A(\mathcal{L}; \underline{t}) & \text{if } \mathcal{L} \text{ is non-empty,} \\ 1 & \text{if } \mathcal{L} \text{ is empty,} \end{cases} \quad (2.69)$$

where $\Phi_1(\mathcal{B}_{\mathcal{L}}; \underline{t})$ is defined by eq. (1.30). Then

$$Q_{\underline{\beta};0}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}) \Big|_{\substack{\lambda_0=0 \\ \lambda=1}} = \prod_{j=1}^{L'} \frac{1 - \left(\prod_{i=1}^L t_i^{l_{ij}} \right)^{-\beta_j}}{1 - \left(\prod_{i=1}^L t_i^{l_{ij}} \right)^{-1}}, \quad (2.70)$$

and the expansion of $Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda})$ in powers of $\underline{t} - 1$ at $\lambda_0 = 0$, $\lambda = 1$ has a form

$$Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}) \Big|_{\substack{\lambda_0=0 \\ \lambda=1}} = \sum_{\underline{m} \geq 0} q_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) (\underline{t} - 1)^{\underline{m}}, \quad (2.71)$$

$$q_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \in \mathbb{Q}[\underline{\beta}], \quad \deg_m q_{\underline{m};n} \leq 2(|\underline{m}| + n) + 1. \quad (2.72)$$

We will prove this proposition in the remainder of this section. However, before we proceed with the detailed calculations of $A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t} | M)$ we want to use Proposition 2.7 in order to derive some properties of the polynomials (2.67).

Proposition 2.8 *The polynomials (2.67) satisfy the following properties*

(1) *If the link $\mathcal{L}$ is empty, then*

$$[\beta_1] q^{\phi(\mathcal{L}' | \underline{\beta})} \Lambda(\underline{\beta}; \underline{\lambda}) \sum_{n=0}^{\infty} Q_{\underline{\beta};n}(\mathcal{L}'; \underline{\lambda}) h^n = J_{\underline{\beta}}(\mathcal{L}'; q | \underline{\lambda}) \quad \text{in } \mathbb{Q}[\underline{\beta}][[h]]. \quad (2.73)$$

(2) *For any j , $1 \leq j \leq L'$*

$$Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}) \Big|_{\beta_j=1} = Q_{\underline{\beta}_{(j)};n}(\mathcal{L} \cup \mathcal{L}'_{(j)}; \underline{t}; \underline{\lambda}). \quad (2.74)$$

(3) *For any i , $1 \leq i \leq L$ and for fixed $\underline{t}_{(i)}$ and $\underline{\lambda}$ such that $Q(\mathcal{L}; \underline{t}; \underline{\lambda})|_{t_i=1} \neq 0$*

$$\begin{aligned} \sum_{\mu=\pm 1} \mu \Phi_2(\mu \alpha_i; \underline{t}_{(i)}; \underline{\lambda}; \underline{\beta}) \sum_{n=0}^k \frac{Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda})}{Q^{2n+1}(\mathcal{L}; \underline{t}; \underline{\lambda})} \Big|_{t_i=q^{\mu \alpha_i}} \\ = \Phi_2(\alpha_i; \underline{t}_{(i)}; \underline{\lambda}; \underline{\beta}) \sum_{n=0}^k \frac{Q_{\alpha_i; \beta; n}(\mathcal{L}_{(i)}, \mathcal{L}_i \cup \mathcal{L}'; \underline{t}_{(i)}; \underline{\lambda})}{Q^{2n+1}(\mathcal{L}_{(i)}; \underline{t}_{(i)}; \underline{\lambda})} + O(h^{k+1}) \end{aligned} \quad (2.75)$$

as $h \rightarrow 0$.

Proof. Equation (2.73) follows from eqs. (2.45) and (2.33). Equation (2.74) follows from eq. (2.46). In order to prove eq. (2.75) we consider the relation eq. (2.63) in the vicinity of $\underline{t} = 1, \underline{\lambda} = 0$ in which a uniform convergence (2.68) holds. Since the substitution $t_i = q^{\mu\alpha_i}$ leaves us within the same vicinity as we set $h \rightarrow 0$, then the uniformity of convergence allows us to apply the limit $M \rightarrow \infty$ to both sides of eq. (2.75). Thus we obtain (2.75) in the vicinity of $\underline{t}_{(i)} = 1, \underline{\lambda} = 0$. Since both sides of eq. (2.75) are rational (and hence analytic) functions, then the relation (2.75) should also hold at any point $\underline{t}_{(i)}, \underline{\lambda}$ such that $Q(\mathcal{L}; \underline{t}; \underline{\lambda})|_{t_i=1} \neq 0$. $\square$

2.2.2 Fixing transitions

We will prove Proposition 2.7 by calculating the limit $\lim_{M \rightarrow \infty} A_{\beta;n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M)$ explicitly. However, before we do this, we have to introduce some notations. Let V_1 and V_2 be two $SU_q(2)$ modules (either finite, or infinite-dimensional) with colors γ_1 and γ_2 and with bases $f_m, m \in \mathcal{S}_1$ and $f_m, m \in \mathcal{S}_2, \mathcal{S}_{1,2} \subset \mathbb{Z}$. For $n \geq 0$ we define two operators $R^{\pm,n}: V_1 \otimes V_2 \rightarrow V_2 \otimes V_1$ such that

$$R^{+,n}(f_{m_1} \otimes f_{m_2}) = \begin{cases} \check{R}_{m_1, m_2}^{m_2+n, m_1-n} f_{m_2+n} \otimes f_{m_1-n} & \text{if } m_2+n \in \mathcal{S}_1, m_1-n \in \mathcal{S}_2, \\ 0 & \text{otherwise,} \end{cases} \quad (2.76)$$

$$R^{-,n}(f_{m_1} \otimes f_{m_2}) = \begin{cases} (\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} f_{m_2-n} \otimes f_{m_1+n} & \text{if } m_2-n \in \mathcal{S}_1, m_1+n \in \mathcal{S}_2, \\ 0 & \text{otherwise.} \end{cases} \quad (2.77)$$

Obviously,

$$\check{R} = \sum_{n \geq 0} R^{+,n}, \quad \check{R}^{-1} = \sum_{n \geq 0} R^{-,n}. \quad (2.78)$$

Let us call the positive $(\mathcal{L}, \mathcal{L}')$ crossings and the negative $(\mathcal{L}', \mathcal{L})$ crossings *the losing crossings* and let us call the negative $(\mathcal{L}, \mathcal{L}')$ and positive $(\mathcal{L}', \mathcal{L})$ crossings *the gaining crossings*. Suppose that the braid $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ has N^- losing crossings, N^+ gaining crossings and N'' crossings of $(\mathcal{L}', \mathcal{L}')$ type and that the crossings within each group are indexed in some order. Then three lists of non-negative numbers $\underline{n}^- = (n_1^-, \dots, n_{N^-}^-)$, $\underline{n}^+ = (n_1^+, \dots, n_{N^+}^+)$ and $\underline{n}'' = (n_1'', \dots, n_{N''}'')$ assign a non-negative number to any crossing other than those of $(\mathcal{L}, \mathcal{L})$

type. After this assignment we construct an operator $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$, $\underline{n} = \underline{n}^-; \underline{n}^+; \underline{n}''$ by performing the following transformation on the expression for $B_{\mathcal{L} \cup \mathcal{L}'; M}$: we replace the operators $\check{R}$ and $\check{R}^{-1}$ at all the crossings except of $(\mathcal{L}, \mathcal{L})$ type by the operators $R^{+,n}$ and $R^{-,n}$ whose numbers n are equal to the numbers assigned by $\underline{n}$ to the corresponding crossings.

If we replace the braiding operator $B_{\mathcal{L} \cup \mathcal{L}'; M}$ by $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$ in eq. (2.47), then we get the trace

$$J_{\underline{\alpha}; \underline{\beta} | \underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) \quad (2.79)$$

$$= [\alpha_*] q^{\phi(\mathcal{L} \cup \mathcal{L}' | \underline{\alpha}, \underline{\beta})} \Lambda(\underline{\alpha}, \underline{\beta}; \underline{\lambda}) \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}} P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda}, *}^{(N)} Q_{q, *}^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}} \quad (2.80)$$

Obviously, the trace (2.79) shares the properties of $J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$ listed in Proposition 2.5 and Corollary 2.6:

Proposition 2.9 *For fixed $\underline{\beta}$ and M there exists a unique set of polynomials*

$$A_{\underline{\beta} | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; \underline{t}; q | M) \in \mathbb{Z}[\underline{\lambda}, \underline{t}, \underline{t}^{-1}, q, q^{-1}], \quad A_{\underline{\beta}; n | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; \underline{t} | M) \in \mathbb{Z}[\underline{\lambda}, \underline{t}, \underline{t}^{-1}], \quad (2.81)$$

$$A_{\underline{\beta} | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; \underline{t}; q | M) = \sum_{n=0}^{\infty} A_{\underline{\beta}; n | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; \underline{t} | M) h^n, \quad (2.82)$$

such that

$$J_{\underline{\alpha}; \underline{\beta} | \underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = [\alpha_*] q^{\phi(\mathcal{L} \cup \mathcal{L}' | \underline{\alpha}, \underline{\beta})} \Lambda(\underline{\alpha}, \underline{\beta}; \underline{\lambda}) A_{\underline{\beta} | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; q^{\underline{\alpha}}; q | M), \quad (2.83)$$

$$A_{\underline{\beta} | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; q^{\underline{\alpha}}; q | M) = \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}} P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda}, *}^{(N)} Q_{q, *}^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}, \quad (2.84)$$

and for fixed $\underline{\alpha}, \underline{\beta}$

$$J_{\underline{\alpha}; \underline{\beta} | \underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = [\alpha_*] q^{\phi(\mathcal{L} \cup \mathcal{L}' | \underline{\alpha}, \underline{\beta})} \Lambda(\underline{\alpha}, \underline{\beta}; \underline{\lambda}) \sum_{n=0}^{\infty} A_{\underline{\beta}; n | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; q^{\underline{\alpha}} | M) h^n. \quad (2.85)$$

Proof. The proof is absolutely similar to that of Proposition 2.5 and Corollary 2.6. $\square$

There is a simple relation between $J_{\underline{\alpha}; \underline{\beta} | \underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$ and $J_{\underline{\alpha}; \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$.

Proposition 2.10

$$J_{\underline{\alpha};\underline{\beta}|\underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = \sum_{\substack{0 \leq \underline{n} \leq 2M \\ |\underline{n}^+| = |\underline{n}^-|}} J_{\underline{\alpha};\underline{\beta}|\underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M), \quad (2.86)$$

$$A_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q | M) = \sum_{\substack{0 \leq \underline{n} \leq 2M \\ |\underline{n}^+| = |\underline{n}^-|}} A_{\underline{\beta}|\underline{n}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q | M), \quad (2.87)$$

$$A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q | M) = \sum_{\substack{0 \leq \underline{n} \leq 2M \\ |\underline{n}^+| = |\underline{n}^-|}} A_{\underline{\beta};n|\underline{n}}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}; q | M). \quad (2.88)$$

Proof. It is easy to see that due to the presence of projection operators P_M in the expression for $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$, its matrix elements are zero unless $\underline{n} \leq 2M$. Therefore in view of eqs.(2.78)

$$J_{\underline{\alpha};\underline{\beta}|\underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M) = \sum_{0 \leq \underline{n} \leq 2M} J_{\underline{\alpha};\underline{\beta}|\underline{n}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M). \quad (2.89)$$

Let $H_{\mathcal{L}'}$ be a sum of $SU_q(2)$ generators H acting on the components of $V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}$ within the tensor product $V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}$. The space $V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}$ is a sum of eigenspaces $(V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty})^{[n]}$ of $\mathcal{L}'$ corresponding to its eigenvalues n

$$V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty} = \bigoplus_{n \in \mathbb{Z}} (V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty})^{[n]} \quad (2.90)$$

It is easy to see that

$$P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda}, * }^{(N)} Q_{q, *}^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}} : \quad (2.91)$$

$$(V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty})^{[n]} \rightarrow (V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty})^{[n+2(|\underline{n}^+| - |\underline{n}^-|)]}.$$

Therefore

$$\text{Tr}_{V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}} P_{\mathcal{L} \cup \mathcal{L}'; M} Q_{\underline{\lambda}, * }^{(N)} Q_{q, *}^{[1]} B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}} = 0, \quad \text{unless } |\underline{n}^+| = |\underline{n}^-|. \quad (2.92)$$

Then eq.(2.86) follows from eq.(2.89). Relations (2.88) follow from eq.(2.86) because of the uniqueness of polynomials A . $\square$

2.2.3 Expansion of matrix elements

Now it's time to expand the matrix elements of the operators $\check{R}$ and $\check{R}^{-1}$ participating in the expression for $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$, as well as the matrix elements of the operator $Q_{q, *}^{[1]}$. Since we

have to study the expansion of the trace $J_{\underline{\alpha}, \underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; q | \underline{\lambda}; M)$ in powers of h while keeping the expressions (2.4) unexpanded then we will need different formulas for expanding the matrix elements (2.19) and (2.20) depending on the type of the crossing that generated them.

Theorem 2.11 *There exist the polynomials*

$$\begin{aligned} T_{j,k}^{(\pm)}(m_1, m_2, n) &\in \mathbb{Q}[m_1, m_2, n], \quad j, k \geq 0, \\ \deg_{m_1} T_{j,k}^{(+)} , \deg_{m_2} T_{j,k}^{(-)} &\leq k, \quad \deg_{m_2} T_{j,k}^{(+)} , \deg_{m_1} T_{j,k}^{(-)} \leq j + k, \\ \deg_{m_1} T_{j,k}^{(\pm)} + \deg_{m_2} T_{j,k}^{(\pm)} + \deg_n T_{j,k}^{(\pm)} &\leq j + 2k, \\ T_{0,0}^{((m_1, m_2, n), \pm)} &= 1, \end{aligned} \tag{2.93}$$

such that for fixed $\gamma_1, \gamma_2, m_1, m_2$, and n

$$\begin{aligned} \check{R}_{m_1, m_2}^{m_2+n, m_1-n} &= q^{(\gamma_1-1)(\gamma_2-1)/4} \binom{m_1}{n} (1 - q^{-\gamma_2})^n q^{-m_2 \gamma_1} \\ &\times \sum_{j=0}^n \left(\frac{\prod_{l=0}^{j-1} (n-l)}{(1 - q^{-\gamma_2})^j} \sum_{k=0}^{\infty} T_{j,k}^{(+)}(m_1, m_2, n) h^{j+k} \right), \end{aligned} \tag{2.94}$$

$$\begin{aligned} (\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} &= q^{-(\gamma_1-1)(\gamma_2-1)/4} \binom{m_2}{n} (1 - q^{\gamma_1})^n q^{m_1 \gamma_2 + n(\gamma_2 - \gamma_1)} \\ &\times \sum_{j=0}^n \left(\frac{\prod_{l=0}^{j-1} (n-l)}{(1 - q^{\gamma_1})^j} \sum_{k=0}^{\infty} T_{j,k}^{(-)}(m_1, m_2, n) h^{j+k} \right). \end{aligned} \tag{2.95}$$

The polynomials $T_{j,k}^{(+)}$ and $T_{j,k}^{(-)}$ are related

$$\sum_{k=0}^{\infty} T_{j,k}^{(-)}(m_1, m_2, n) h^{j+k} = \sum_{k=0}^{\infty} T_{j,k}^{(+)}(m_2, m_1, n) \left(-\frac{h}{1+h} \right)^{j+k}. \tag{2.96}$$

This theorem essentially says that there are at most two powers of m_1, m_2 or n for each power of h in the expansion of the matrix elements. This is easy to see by inspecting eqs. (2.19) and (2.20). The only non-trivial part of the theorem is that j powers of $(1 - q^{\pm \gamma_{1,2}})$ in denominators of eqs. (2.94), (2.95) are accompanied by the factor $\prod_{l=0}^{j-1} (n-l)$.

Proof of Theorem 2.11. This theorem is a simple corollary of Lemma 3.1 of [8]. $\square$

We present explicit formulas for some polynomials $T_{j,k}^{(\pm)}$ in Appendix B.

Corollary 2.12 *There are three other ways to expand the matrix elements $\check{R}_{m_1, m_2}^{m_2+n, m_1-n}$ and $(\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n}$.*

(1) *There exist the polynomials*

$$\begin{aligned} T_{n,k}^{(1,\pm)}(m_1, m_2, \gamma) &\in \mathbb{Q}[m_1, m_2, \gamma], \\ \deg_{m_1} T_{n,k}^{(1,\pm)} &\leq k, \quad \deg_{\gamma} T_{n,k}^{(1,\pm)} + \deg_{m_2} T_{n,k}^{(1,\pm)} \leq n+k, \\ T_{0,0}^{(1,\pm)}(m_1, m_2, \gamma) &= 1, \end{aligned} \quad (2.97)$$

such that

$$\check{R}_{m_1, m_2}^{m_2+n, m_1-n} = q^{(\gamma_1-1)(\gamma_2-1)/4} \binom{m_1}{n} q^{-m_2\gamma_1} \sum_{k=0}^{\infty} T_{n,k}^{(1,+)}(m_1, m_2, \gamma_2) h^{n+k}, \quad (2.98)$$

$$(\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} = q^{-(\gamma_1-1)(\gamma_2-1)/4} \binom{m_2}{n} q^{(m_1+n)\gamma_2} \sum_{k=0}^{\infty} T_{n,k}^{(1,-)}(m_1, m_2, \gamma_1) h^{n+k} \quad (2.99)$$

(2) *There exist the polynomials*

$$\begin{aligned} T_{n,k}^{(2,\pm)}(m_1, m_2, \gamma, t) &\in \mathbb{Q}[m_1, m_2, \gamma, t, t^{-1}], \\ \deg_{m_2} T_{n,k}^{(2,+)} &\leq k, \quad \deg_{m_1} T_{n,k}^{(2,-)} \leq k, \\ T_{0,0}^{(2,\pm)}(m_1, m_2, \gamma, t) &= 1, \end{aligned} \quad (2.100)$$

such that there is a bound on the powers of m and γ in the coefficients of the expansion of the polynomials in powers of $t-1$

$$T_{n,k}^{(2,\pm)}(m_1, m_2, \gamma, t) = \sum_{l=0}^{\infty} T_{n,k,l}^{(2,\pm)}(m_1, m_2, \gamma) (t-1)^l,$$

$$T_{n,k,l}^{(2,\pm)}(m_1, m_2, \gamma) \in \mathbb{Q}[m_1, m_2, \gamma],$$

$$\deg_{m_1} T_{n,k,l}^{(2,+)} + \deg_{\gamma} T_{n,k,l}^{(2,+)} \leq k+l, \quad (2.101)$$

$$\deg_{m_2} T_{n,k,l}^{(2,-)} + \deg_{\gamma} T_{n,k,l}^{(2,-)} \leq k+l, \quad (2.102)$$

and for fixed $\gamma_1, \gamma_2, m_1, m_2$, and n

$$\check{R}_{m_1, m_2}^{m_2+n, m_1-n} = q^{(\gamma_1-1)(\gamma_2-1)/4} \sum_{k=0}^{\infty} T_{n,k}^{(2,+)}(m_1, m_2, \gamma_1, q^{\gamma_2}) h^{n+k}, \quad (2.103)$$

$$(\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} = q^{-(\gamma_1-1)(\gamma_2-1)/4} \sum_{k=0}^{\infty} T_{n,k}^{(2,-)}(m_1, m_2, \gamma_2, q^{\gamma_1}) h^{n+k}. \quad (2.104)$$

(3) *There exist polynomials*

$$\begin{aligned} T_{n,k}^{(3,\pm)}(\gamma_1, \gamma_2, m_1, m_2) &\in \mathbb{Q}[\gamma_1, \gamma_2, m_1, m_2], \\ \deg_{m_1} T_{n,k}^{(2,\pm)} + \deg_{\gamma_1} T_{n,k}^{(2,\pm)} &\leq k, \quad \deg_{m_2} T_{n,k}^{(2,\pm)} + \deg_{\gamma_2} T_{n,k}^{(2,\pm)} \leq k, \quad (2.105) \\ T_{0,0}^{(3,\pm)}(\gamma_1, \gamma_2, m_1, m_2) &= 1, \end{aligned}$$

such that for fixed $\gamma_1, \gamma_2, m_1, m_2$, and n

$$\check{R}_{m_1, m_2}^{m_2+n, m_1-n} = q^{(\gamma_1-1)(\gamma_2-1)/4} \sum_{k=0}^{\infty} T_{n,k}^{(3,+)}(\gamma_1, \gamma_2, m_1, m_2) h^{n+k}, \quad (2.106)$$

$$(\check{R}^{-1})_{m_1, m_2}^{m_2-n, m_1+n} = q^{-(\gamma_1-1)(\gamma_2-1)/4} \sum_{k=0}^{\infty} T_{n,k}^{(3,-)}(\gamma_1, \gamma_2, m_1, m_2) h^{n+k}. \quad (2.107)$$

Proof. One simply has to expand eqs.(2.94) and (2.95) in the corresponding parameters. We leave the details to the reader. $\square$

Finally, we expand the matrix elements of the operators $q^{\hat{m}}$ which comprise the operator $Q_{q,*}^{[1]}$. The matrix elements of $q^{\hat{m}}$ acting on V_γ are all diagonal and

$$(q^{\hat{m}})_m^m = q^m = \sum_{k=0}^{\infty} \binom{m}{k} h^k. \quad (2.108)$$

(note that $\binom{m}{k} = 0$ for $k > m$ if we treat $\binom{m}{k}$ as a polynomial in m of degree k).

Proposition 2.13 *Suppose that we make the following substitution in the expression for the regularized braiding matrix $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$: we use expressions (2.94), (2.95) for the matrix elements of $\check{R}$ and $\check{R}^{-1}$ at $(\mathcal{L}, \mathcal{L})$ crossings, expressions (2.98), (2.99) at losing crossings, expressions (2.103), (2.104) at gaining crossings and expressions (2.106), (2.107) at $(\mathcal{L}', \mathcal{L}')$ crossings. Suppose that we also use the expression (2.108) for the matrix elements of the operator $Q_q^{[1]}$. Then after this substitution the trace (2.79) transforms into the series (2.85).*

Proof. The presence of the projection operator $P_{\mathcal{L} \cup \mathcal{L}'; M}$ in the trace (2.79) guarantees that only a finite number of matrix elements participating in that trace is non-zero. Therefore the non-zero products of expanded matrix elements yield a series of the form (2.85) with $A_{\underline{\beta}; \underline{n} | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; \underline{t} | M) \in \mathbb{Q}[\underline{\lambda}, \underline{t}, \underline{t}^{-1}]$, and this series converges uniformly in a vicinity of $h = 0$. Now the claim of the proposition follows from the uniqueness of the expansion (2.85). $\square$

Corollary 2.14 *The bounds in the sum of eq. (2.88) can be made M -independent*

$$A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t}|M) = \sum_{\substack{\underline{n} \geq 0 \\ |\underline{n}^+| = |\underline{n}^-| \\ |\underline{n}^-| + |\underline{n}''| \leq n}} A_{\underline{\beta};n|\underline{n}}(\mathcal{L}, \mathcal{L}'\underline{\lambda}; \underline{t}|M). \quad (2.109)$$

Proof. Expansions (2.98), (2.99) of the matrix elements which appear at losing crossings and expansions (2.106), (2.107) of the matrix elements which appear at the $(\mathcal{L}', \mathcal{L}')$ crossings all start at h^n . Since, according to Proposition (2.13), these expansions lead to eq. (2.85), then we conclude that

$$A_{\underline{\beta}|\underline{n}}(\mathcal{L}, \mathcal{L}'\underline{\lambda}; \underline{t}; q|M) = \sum_{n \geq |\underline{n}^-| + |\underline{n}''|} A_{\underline{\beta};n|\underline{n}}(\mathcal{L}, \mathcal{L}'\underline{\lambda}; \underline{t}|M) h^n. \quad (2.110)$$

If we substitute this expansion into eq. (2.87) and compare the result with eq. (2.65), then we come to eq. (2.109). $\square$

2.2.4 Parametrized matrices

Now we are going to use a special trick in order to calculate the trace of the expanded matrices. This trick is a usual method of calculating perturbative expansions in quantum field theory. Its application is mathematically rigorous in the finite-dimensional calculations of this paper. The idea is that in order to calculate a trace of a complicated matrix one may introduce a simpler matrix which depends on some artificial parameters and which satisfies two properties: its trace is easy to calculate and the complicated matrix can be expressed as a combination of derivatives of the parametrized matrix over its parameters.

The first step in implementing our strategy is to construct a matrix $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}[\underline{t}; \underline{\epsilon}]$ depending on a list of parameters $\underline{t}; \underline{\epsilon}$, together with a differential operator $D_{\underline{n}}(\underline{t}; \underline{\beta}; \partial_{\underline{\epsilon}}; h)$ which is a power series in h whose coefficients are polynomials in the derivatives $\partial_{\underline{\epsilon}}$. We will construct $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}[\underline{t}; \underline{\epsilon}]$ in such a way that the matrix elements of $D_{\underline{n}}(\underline{t}; \underline{\beta}; \partial_{\underline{\epsilon}}; h) B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}[\underline{t}; \underline{\epsilon}] \Big|_{\substack{\underline{t}=q\alpha \\ \underline{\epsilon}=0}}$ are equal to those of the matrix $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$ expanded as described in Proposition 2.13. In fact, we will have to consider an extra $SU_q(2)$ module $V_{1,\infty}$, so that our matrix $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}[\underline{t}; \underline{\epsilon}]$ will act

on a tensor product $V_{1,\infty} \otimes V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha},\infty} \otimes V_{\mathcal{B}_{\mathcal{L}'};\underline{\beta},\pm\infty}$ and the expansion of $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}$ will be reproduced by an operator which is a partial trace $D_{\underline{n}}(\underline{t}; \underline{\beta}; \partial_{\underline{\epsilon}}; h) \text{Tr}_{V_{1,\infty}} \lambda_*^{-H} B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}[\underline{t}; \underline{\epsilon}] \Big|_{\substack{\underline{t}=q\alpha \\ \lambda_*, \underline{\epsilon}=0}}.$

The basis of our construction is the matrix $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}$ itself. We will present every $\check{R}, \check{R}^{-1}$, $R^{\pm,n}$ participating in the expression for $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}$ as $D(\partial_{\underline{\epsilon}})R[\underline{\epsilon}]$, where $D(\partial_{\underline{\epsilon}})$ is a power series in h with coefficients which are polynomials in the derivatives over some parameters $\underline{\epsilon}$, and $R[\underline{\epsilon}]$ are operators depending on parameters $\underline{\epsilon}$. These expressions will match the expansion formulas of Theorem 2.11 and Corollary 2.12. Then we will replace the original matrices of $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}$ by $R[\underline{\epsilon}]$ thus producing $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}[\underline{t}; \underline{\epsilon}]$, while the operator $D_{\underline{n}}(\underline{t}; \underline{\beta}; \partial_{\underline{\epsilon}}; h)$ will be the product of individual operators $D(\partial_{\underline{\epsilon}})$.

We begin by defining a parametrized matrix $R^{\pm}[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4]$, $\underline{\epsilon} = (e_1, e_2, e_3, e_4)$ which acts on a tensor product of two Verma modules $V_{\alpha_1,\infty} \otimes V_{\alpha_2,\infty}$:

$$R^+[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4](f_{m_1} \otimes f_{m_2}) \quad (2.111)$$

$$= \sum_{0 \leq n \leq m_1} (R^+[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4])_{m_1, m_2}^{m_2+n, m_1-n} f_{m_2+n} \otimes f_{m_1-n},$$

$$R^-[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4](f_{m_1} \otimes f_{m_2}) \quad (2.112)$$

$$= \sum_{0 \leq n \leq m_2} (R^-[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4])_{m_1, m_2}^{m_2-n, m_1+n} f_{m_2-n} \otimes f_{m_1+n},$$

where

$$(R^+[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4])_{m_1, m_2}^{m_2+n, m_1-n} = \binom{m_1}{n} (1 - t_2^{-1} + \epsilon_4)^n t_1^{-m_2} e^{m_1 \epsilon_1 + m_2 \epsilon_2 + n \epsilon_3}, \quad (2.113)$$

$$(R^-[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4])_{m_1, m_2}^{m_2-n, m_1+n} = \binom{m_2}{n} (1 - t_1 + \epsilon_4)^n t_1^{-n} t_2^{m_1+n} e^{m_1 \epsilon_1 + m_2 \epsilon_2 + n \epsilon_3}. \quad (2.114)$$

The matrix elements (2.113), (2.114) are related to those of (2.19), (2.20), because according to eqs. (2.94), (2.95)

$$\check{R}_{m_1, m_2}^{m_2+n, m_1-n} = q^{(\alpha_1-1)(\alpha_2-1)/4} \quad (2.115)$$

$$\times \left(\sum_{j,k=0}^{\infty} T_{j,k}^{(+)}(\partial_{\epsilon_1}, \partial_{\epsilon_2}, \partial_{\epsilon_3}) \partial_{\epsilon_4}^j h^{j+k} \right) (R^+[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4])_{m_1, m_2}^{m_2+n, m_1-n} \Big|_{\substack{t_{1,2}=q^{\alpha_{1,2}} \\ \epsilon=0}},$$

$$(\check{R}^{-1})_{m_1, m_2}^{m_2+n, m_1-n} = q^{-(\alpha_1-1)(\alpha_2-1)/4} \quad (2.116)$$

$$\times \left(\sum_{j,k=0}^{\infty} T_{j,k}^{(-)}(\partial_{\epsilon_1}, \partial_{\epsilon_2}, \partial_{\epsilon_3}) \partial_{\epsilon_4}^j h^{j+k} \right) (R^-[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4])_{m_1, m_2}^{m_2-n, m_1+n} \Big|_{\substack{t_{1,2}=q^{\alpha_{1,2}} \\ \epsilon=0}}.$$

Note that we could replace the sums $\sum_{j=0}^n$ of eqs. (2.94), (2.95) with the sums $\sum_{j=0}^\infty$, because the terms with $j > n$ in eqs. (2.115), (2.116) are zero due to the fact that if $j > n$, then $\partial_{\epsilon_4}^j (1 - t_{1,2}^{\mp 1} + \epsilon_4)^n = 0$. Following our strategy, we replace the operators $\check{R}, \check{R}^{-1}$ which appear in the expression for $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$ at the $(\mathcal{L}, \mathcal{L})$ crossings, with operators $R^\pm[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4]$, while the product of operators $\sum_{j=0}^\infty \left(\partial_{\epsilon_4}^j \sum_{k=0}^\infty T_{j,k}^{(+)}(\partial_{\epsilon_1}, \partial_{\epsilon_2}, \partial_{\epsilon_3}) h^{j+k} \right)$ coming from all $(\mathcal{L}, \mathcal{L})$ crossings is the first factor of $D_{\underline{n}}(\underline{t}; \underline{\beta}; \partial_{\underline{\epsilon}}; h)$.

Matrix elements (2.98), (2.99) and (2.106), (2.107) require a more elaborate treatment. We define the operators $R^{(\pm)}[\epsilon_1^\pm, \epsilon_2^\pm]$ which act on $V_{1,\infty} \otimes V_{\alpha,\infty}$ according to the formula

$$R^{(+)}[\epsilon_1, \epsilon_2](f_{m_0} \otimes f_{m_1}) = \sum_{0 \leq n \leq m_0} (R^{(+)}[\epsilon_1, \epsilon_2])_{m_0, m_1}^{m_0-n, m_1+n} f_{m_0-n} \otimes f_{m_1+n}, \quad (2.117)$$

$$R^{(-)}[\epsilon_1, \epsilon_2](f_{m_0} \otimes f_{m_1}) = \sum_{0 \leq n \leq m_1} (R^{(-)}[\epsilon_1, \epsilon_2])_{m_0, m_1}^{m_0+n, m_1-n} f_{m_0+n} \otimes f_{m_1-n}, \quad (2.118)$$

where

$$(R^{(+)}[\epsilon_1, \epsilon_2])_{m_0, m_1}^{m_0-n, m_1+n} = \binom{m_0}{n} e^{\epsilon_1 m_1} (\epsilon_2)^n, \quad (2.119)$$

$$(R^{(-)}[\epsilon_1, \epsilon_2])_{m_0, m_1}^{m_0+n, m_1-n} = \binom{m_1}{n} e^{\epsilon_1 m_1} (\epsilon_2)^n. \quad (2.120)$$

We also define another pair of operators depending on parameters t_1, t_2 which act on $V_{\gamma, \pm\infty}$

$$O_n^\pm[t_1, t_2](f_m) = t_1^m t_2^n f_{m \pm n}, \quad (2.121)$$

and an operator C_n which acts on the space $V_{1,\infty}$ as

$$C_n(f_m) = \frac{f_m}{\binom{m}{n}}. \quad (2.122)$$

Now we have to consider four types of mixed crossings.

1. Losing crossings.

(a) Positive $(\mathcal{L}, \mathcal{L}')$ crossing. Consider an operator

$$R_n^{(-,+)}[t_1; \epsilon_1, \epsilon_2, \epsilon'] : V_{1,\infty} \otimes V_{\alpha_1,\infty} \otimes V_{\beta_2, \pm\infty} \rightarrow V_{1,\infty} \otimes V_{\beta_2, \pm\infty} \otimes V_{\alpha_1,\infty}, \quad (2.123)$$

$$R_n^{(-,+)}[t_1; \epsilon_1, \epsilon_2, \epsilon'] = (I \otimes P) (R^{(-)}[\epsilon_1, \epsilon_2] \otimes O_n^+[e^{\epsilon'} t_1^{-1}, 1]). \quad (2.124)$$

It is easy to see that according to eq.(2.98),

$$\delta_{m_0+n}^{m'_0} (R^{-,n})_{m_1,m_2}^{m'_2,m'_1} = q^{(\alpha_1-1)(\beta_2-1)/4} \quad (2.125)$$

$$\times \left(\frac{1}{n!} \partial_{\epsilon_2}^n \sum_{k=0}^{\infty} T_{n,k}^{(1,+)}(\partial_{\epsilon_1}, \partial_{\epsilon'}, \beta_2) h^{n+k} \right) (R_n^{(-,+)}[t_1; \epsilon_1, \epsilon_2, \epsilon'])_{m_0,m_1,m_2}^{m'_0,m'_2,m'_1} \Big|_{\substack{t_1=q^{\alpha_1} \\ \epsilon_1,\epsilon_2,\epsilon'=0}},$$

where $\delta_{m_0+n}^{m'_0}$ is the Kronecker symbol.

(b) Negative $(\mathcal{L}', \mathcal{L})$ crossing. Consider an operator

$$R_n^{(-,-)}[t_2; \epsilon_1, \epsilon_2, \epsilon'] : V_{1,\infty} \otimes V_{\beta_1,\pm\infty} \otimes V_{\alpha_2,\infty} \rightarrow V_{1,\infty} \otimes V_{\alpha_2,\infty} \otimes V_{\beta_1,\pm\infty}, \quad (2.126)$$

$$R_n^{(-,-)}[t_2; \epsilon_1, \epsilon_2, \epsilon'] = (R^{(-)}[\epsilon_1, \epsilon_2] \otimes O_n^+[e^{\epsilon'} t_2, t_2]) (I \otimes P).$$

According to eq.(2.98),

$$\delta_{m_0+n}^{m'_0} (R^{-,n})_{m_1,m_2}^{m'_2,m'_1} = q^{-(\beta_1-1)(\alpha_2-1)/4} \quad (2.127)$$

$$\times \left(\frac{1}{n!} \partial_{\epsilon_2}^n \sum_{k=0}^{\infty} T_{n,k}^{(1,+)}(\partial_{\epsilon'}, \partial_{\epsilon_1}, \beta_1) h^{n+k} \right) (R_n^{(-,-)}[t_2; \epsilon_1, \epsilon_2, \epsilon'])_{m_0,m_1,m_2}^{m'_0,m'_2,m'_1} \Big|_{\substack{t_2=q^{\alpha_2} \\ \epsilon_1,\epsilon_2,\epsilon'=0}}.$$

2. Gaining crossings.

(a) Positive $(\mathcal{L}', \mathcal{L})$ crossing. Consider an operator

$$R_n^{(+,+)}[t_2; \epsilon_1, \epsilon_2, \epsilon'] : V_{1,\infty} \otimes V_{\beta_1,\pm\infty} \otimes V_{\alpha_2,\infty} \rightarrow V_{1,\infty} \otimes V_{\alpha_2,\infty} \otimes V_{\beta_1,\pm\infty}, \quad (2.128)$$

$$R_n^{(+,+)}[t_2; \epsilon_1, \epsilon_2, \epsilon'] = (R^{(+)}[\epsilon_1, \epsilon_2] \otimes O_n^-[e^{\epsilon'}, 1]) (C_n \otimes P). \quad (2.129)$$

According to eq.(2.103),

$$\delta_{m_0-n}^{m'_0} (R^{+,n})_{m_1,m_2}^{m'_2,m'_1} = q^{(\gamma_1-1)(\gamma_2-1)/4} \beta \alpha \quad (2.130)$$

$$\times \left(\frac{1}{n!} \partial_{\epsilon_2}^n \sum_{k=0}^{\infty} T_{n,k}^{(2,+)}(\partial_{\epsilon'}, \partial_{\epsilon_1}, \beta_1, t_2) h^{n+k} \right) (R_n^{(+,+)}[t_2; \epsilon_1, \epsilon_2, \epsilon'])_{m_0,m_1,m_2}^{m'_0,m'_2,m'_1} \Big|_{\substack{t_2=q^{\alpha_2} \\ \epsilon_1,\epsilon_2,\epsilon'=0}}.$$

(b) Negative $(\mathcal{L}, \mathcal{L}')$ crossing. Consider an operator

$$R_n^{(+,-)}[t_1; \epsilon_1, \epsilon_2, \epsilon'] : V_{1,\infty} \otimes V_{\alpha_1,\infty} \otimes V_{\beta_2,\pm\infty} \rightarrow V_{1,\infty} \otimes V_{\beta_2,\pm\infty} \otimes V_{\alpha_1,\infty}, \quad (2.131)$$

$$R_n^{(+,-)}[t_1; \epsilon_1, \epsilon_2, \epsilon'] = (I \otimes P) (R^{(+)}[\epsilon_1, \epsilon_2] \otimes O_n^-[e^{\epsilon'}, 1]) (C_n \otimes I^{\otimes 2}). \quad (2.132)$$

According to eq.(2.104),

$$\delta_{m_0-n}^{m'_0} (R^{+,n})_{m_1,m_2}^{m'_2,m'_1} = q^{-(\gamma_1-1)(\gamma_2-1)/4} \alpha \beta \quad (2.133)$$

$$\times \left(\frac{1}{n!} \partial_{\epsilon_2}^n \sum_{k=0}^{\infty} T_{n,k}^{(2,+)}(\partial_{\epsilon_1}, \partial_{\epsilon'}, \beta_2, t_1) h^{n+k} \right) (R_n^{(+,+)}[t_1; \epsilon_1, \epsilon_2, \epsilon'])_{m_0,m_1,m_2}^{m'_0,m'_2,m'_1} \Big|_{\substack{t_1=q^{\alpha_1} \\ \epsilon_1, \epsilon_2, \epsilon'=0}}.$$

Finally, we turn to $(\mathcal{L}', \mathcal{L}')$ crossings. We introduce two operators

$$R''_{n,\pm}[\epsilon'_1, \epsilon'_2]: V_{\beta_1, \pm\infty} \otimes V_{\beta_2, \pm\infty} \rightarrow V_{\beta_2, \pm\infty} \otimes V_{\beta_1, \pm\infty}, \quad (2.134)$$

$$R''_{n,\pm}[\epsilon'_1, \epsilon'_2] = P(O_n^{\mp}[e^{\epsilon'_1}, 1] \otimes O_n^{\pm}[e^{\epsilon'_2}, 1]), \quad (2.135)$$

so that according to eqs.(2.106), (2.107),

$$(R^{\pm,n})_{m_1,m_2}^{m'_2,m'_1} = q^{\pm(\beta_1-1)(\beta_2-1)/4} \left(T_{n,k}^{(3,\pm)}(\beta_1, \beta_2, \partial_{\epsilon'_1}, \partial_{\epsilon'_2}) h^{n+k} \right) (R''_{n,\pm}[\epsilon'_1, \epsilon'_2])_{m_1,m_2}^{m'_2,m'_1} \Big|_{\epsilon'_1, \epsilon'_2=0} \quad (2.136)$$

Now, according to our strategy, we take the expression for $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}$ and replace $\check{R}$ and $\check{R}^{-1}$ matrices at $(\mathcal{L}, \mathcal{L})$ crossings with the matrices (2.111), (2.112), replace $R^{\pm,n}$ matrices at losing crossings with matrices (2.124), (2.127), replace $R^{\pm,n}$ matrices at gaining crossings with matrices (2.129), (2.132) and replace $R^{\pm,n}$ matrices at $(\mathcal{L}', \mathcal{L}')$ crossings with matrices (2.135). The result is the parametrized matrix $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}^{(C)}[t; \underline{\epsilon}]$ acting on $V_{1,\infty} \otimes V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm\infty}$, where $\underline{\epsilon} = (\underline{\epsilon}; \underline{\epsilon}')$, $\underline{\epsilon}$ and $\underline{\epsilon}'$ being all the parameters ϵ and ϵ' from the individual parametrized matrices. We assume that the matrices (2.124), (2.127), (2.129) and (2.132) act on the space $V_{1,\infty}$ in the following order: losing crossing matrices precede gaining crossing matrices and among themselves those matrices are ordered according to the indexing of losing and gaining crossings.

The operator $B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}^{(C)}[t; \underline{\epsilon}]$ is a tensor product of operators

$$B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}^{(C)}[t; \underline{\epsilon}] = B_{\mathcal{L}; \underline{n}}^{(C)}[t; \underline{\epsilon}] \otimes \bigotimes_{j=1}^{L'} B_{\mathcal{L}'_j; M; \underline{n}}[t; \epsilon'_j], \quad (2.137)$$

$$B_{\mathcal{L}; \underline{n}}^{(C)}[t; \underline{\epsilon}]: V_{1,\infty} \otimes V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \rightarrow V_{1,\infty} \otimes V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty}, \quad B_{\mathcal{L}'_j; M; \underline{n}}[t; \epsilon'_j]: V_{\mathcal{B}_{\mathcal{L}'_j}; \beta_j, \pm\infty} \rightarrow V_{\mathcal{B}_{\mathcal{L}'_j}; \beta_j, \pm\infty}$$

(note that $B_{\mathcal{L}; \underline{n}}^{(C)}[t; \underline{\epsilon}]$ does not depend on M). To construct the operator $B_{\mathcal{L}; \underline{n}}^{(C)}[t; \underline{\epsilon}]$, we place the operators R^+ at positive elementary crossings of $\mathcal{L}$ and the operators R^- at negative elementary crossings of $\mathcal{L}$. We also place the operators $R^{(-)}$ at losing crossings and $R^{(+)} (I \otimes C_n)$

at gaining crossings. To construct the operators $B_{\mathcal{L}'_j;M;\underline{n}}[\underline{t};\underline{\epsilon}']$ we simply place the appropriate operators $O_n^\pm$ at every place where a strand of $\mathcal{L}_j$ crosses some other strand.

We construct the differential operator $D_{\underline{n}}(\underline{t};\underline{\beta};\partial_{\underline{\epsilon}};h)$ simultaneously with $B_{\mathcal{L}'_j;M;\underline{n}}[\underline{t};\underline{\epsilon}']$ by taking the products of differential operators which accompany individual parametrized matrices.

Proposition 2.15 *The operator $D_{\underline{n}}(\underline{t};\underline{\beta};\partial_{\underline{\epsilon}};h)$ has the following form:*

$$D_{\underline{n}}(\underline{t};\underline{\beta};\partial_{\underline{\epsilon}};h) = \sum_{n=0}^{\infty} D_{\underline{n};n}(\underline{t};\underline{\beta};\partial_{\underline{\epsilon}}) h^n, \quad D_{\underline{n};n}(\underline{t};\underline{\beta};\underline{\epsilon}) \in \mathbb{Q}[\underline{t}, \underline{t}^{-1}, \underline{\beta}, \underline{\epsilon}], \quad (2.138)$$

$$D_{\underline{0};0}(\underline{t};\underline{\beta};\underline{\epsilon}) = 1. \quad (2.139)$$

The expansion of the polynomials $D_{\underline{n};n}(\underline{t};\underline{\beta};\underline{\epsilon})$ in powers of $\underline{t} - 1$ has the form

$$D_{\underline{n};n}(\underline{t};\underline{\beta};\underline{\epsilon}) = \sum_{\underline{m} \geq 0} d_{\underline{n};n;\underline{m}}(\underline{\beta};\underline{\epsilon}) (\underline{t} - 1)^{\underline{m}}, \quad d_{\underline{n};n;\underline{m}}(\underline{\beta};\underline{\epsilon}) \in \mathbb{Q}[\underline{\beta}, \underline{\epsilon}], \quad (2.140)$$

$$\deg_{\underline{\beta}} d_{\underline{n};n;\underline{m}} + \deg_{\underline{\epsilon}} d_{\underline{n};n;\underline{m}} \leq |\underline{m}| + n.$$

Proof. Equation (2.139) follows from the similar relations in eqs. (2.93), (2.97), (2.100) and (2.105). Equations (2.138) and (2.140) follow easily from Theorem 2.11 and Corollary 2.12. $\square$

We chose the matrix elements of gaining and losing crossing operators $R_n^{(\pm)}$ in such a way that if we take any diagonal element of the operator $D_{\underline{n}}(\underline{t};\underline{\beta};\partial_{\underline{\epsilon}};h)B_{\mathcal{L}'_j;M;\underline{n}}[\underline{t};\underline{\epsilon}']$ at $|\underline{n}^+| = |\underline{n}^-|$ in the space $V_{1,\infty}$, then, according to Proposition 2.13, the resulting operator acting on $V_{\mathcal{B}_{\mathcal{L}};\underline{\alpha},\infty} \otimes V_{\mathcal{B}_{\mathcal{L}'};\underline{\beta},\pm\infty}$ coincides with the expansion of $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}$, as described in that proposition. In particular, if we take a diagonal element corresponding to f_0 , then

$$B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}} = D_{\underline{n}}(\underline{t};\underline{\beta};\partial_{\underline{\epsilon}};h) \text{Tr}_{V_{1,\infty}}(P_M \lambda_*^{2\hat{m}} \otimes I^{\otimes N}) B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}^{(C)}[\underline{t};\underline{\epsilon}] \Big|_{\substack{\lambda_* = 0 \\ \underline{t} = q\hat{\underline{\alpha}} \\ \underline{\epsilon} = 0}} \quad (2.141)$$

(here we used an obvious fact that $P_M f_0 = f_0$).

Let us remove all operators C_n from the expressions for $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}^{(C)}[\underline{t};\underline{\epsilon}]$ and $B_{\mathcal{L};\underline{n}}^{(C)}[\underline{t};\underline{\epsilon}]$. We denote the resulting operators as $B_{\mathcal{L} \cup \mathcal{L}';M;\underline{n}}[\underline{t};\underline{\epsilon}]$ and $B_{\mathcal{L}}[\underline{t};\underline{\epsilon}]$ (note that $B_{\mathcal{L}}[\underline{t};\underline{\epsilon}]$ no longer depends on $\underline{n}$). Since the change in the index m_0 of the vector $f_{m_0} \in V_{1,\infty}$, as it passes through

the operators (2.125), (2.127), (2.130) and (2.133), is fixed by the Kronecker symbols, then the removal of the operators C_n from the *r.h.s.* of eq.(2.141) can be compensated by dividing it by the product of factorials coming from eq.(2.122)

$$B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}} = F(\underline{n}^+) D_{\underline{n}}(\underline{t}; \underline{\beta}; \partial_{\underline{\epsilon}}; h) \text{Tr}_{V_{1, \infty}}(I^{\otimes N} \otimes P_M \lambda_*^{2\hat{m}}) B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}^{(C)}[\underline{t}; \underline{\epsilon}] \Big|_{\substack{\lambda_* = 0 \\ \underline{t} = q \frac{\underline{\alpha}}{\underline{\epsilon}} \\ \underline{\epsilon} = 0}}, \quad (2.142)$$

$$F(\underline{n}^+) = \left(\sum_{i=1}^{N^+} \binom{|\underline{n}^+| - \sum_{j=1}^{i-1} n_j^+}{n_i^+} \right)^{-1}. \quad (2.143)$$

It remains to ‘parametrize’ the expansion (2.108) of the matrix elements of the operator $Q_{\underline{\lambda}, *}$. We introduce the numbers $C_{k, n} \in \mathbb{Q}$ according to the formula

$$\frac{1}{n!} \left((1+h)^{-1/2} - 1 \right)^n = \sum_{k \geq n} C_{k, n} h^k. \quad (2.144)$$

Note that

$$C_{0, 0} = 1 \quad (2.145)$$

Since

$$\begin{aligned} \lambda^{2m} q^{-m} &= \left(\lambda + \left((1+h)^{-1/2} - 1 \right) \lambda \right)^{2m} \\ &= \left(\sum_{n=0}^{\infty} \frac{\left((1+h)^{-1/2} - 1 \right)^n}{n!} (\lambda \partial_{\lambda})^n \right) \lambda^{2m} \\ &= \left(\sum_{\substack{k \geq 0 \\ 0 \leq n \leq k}} C_{k, n} h^k (\lambda \partial_{\lambda})^n \right) \lambda^{2m}, \end{aligned} \quad (2.146)$$

then the *r.h.s.* of the following equation represents the expansion of the matrix elements of $Q_{q, *}^{[1]}$ described by eq.(2.108)

$$Q_{\underline{\lambda}, *}^{(N)} Q_{q, *}^{[1]} = D(\lambda; \partial_{\lambda}; h) Q_{\underline{\lambda}, *}^{(N)}, \quad (2.147)$$

where

$$D(\lambda; \partial_{\lambda}; h) = \sum_{\substack{k \geq 0 \\ 0 \leq n \leq k}} C_{k, n} h^k \lambda^n \partial_{\lambda}^n. \quad (2.148)$$

Consider a product of operators

$$D_{\underline{n}}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda}; h) = D(\lambda; \partial_{\lambda}; h) D_{\underline{n}}(\underline{t}; \underline{\beta}; \partial_{\underline{\epsilon}}; h) \quad (2.149)$$

Proposition 2.16 *The operator $D_{\underline{n}}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda}; h)$ is a power series in h*

$$D_{\underline{n}}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda}; h) = \sum_{n=0}^{\infty} D_{\underline{n};n}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda}) h^n, \quad D_{\underline{n};n}(\underline{t}; \lambda; \underline{\beta}; \underline{\epsilon}; x) \in \mathbb{Q}[\underline{t}, \underline{t}^{-1}, \lambda, \underline{\beta}, \underline{\epsilon}, x],$$

$$D_{\underline{0},0}(\underline{t}; \lambda; \underline{\beta}; \underline{\epsilon}; x) = 1, \quad (2.150)$$

$$\deg_{\underline{\epsilon}} D_{\underline{n};n} + \deg_x D_{\underline{n};n} \leq 2n, \quad (2.151)$$

$$D_{\underline{n};n}(\underline{t}; \underline{\lambda}; \underline{\beta}; \underline{\epsilon}; x) = \sum_{\underline{m} \geq 0} d_{\underline{m},\underline{n};n}(\underline{\lambda}; \underline{\beta}; \underline{\epsilon}; x) (\underline{t} - 1)^{\underline{m}}, \quad (2.152)$$

$$\deg_x d_{\underline{m},\underline{n};n} + \deg_{\underline{\beta}} d_{\underline{m},\underline{n};n} + \deg_{\underline{\epsilon}'} d_{\underline{m},\underline{n};n} \leq 2n + |\underline{m}|. \quad (2.153)$$

and its action on the trace of the operator $Q_{\underline{\lambda},*}^{(N)} B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}[\underline{t}; \underline{\epsilon}]$ yields the expansion (2.83), so that

$$A_{\underline{\beta};n|\underline{n}}(\mathcal{L}, \mathcal{L}'\underline{\lambda}; \underline{t}|M) = F(\underline{n}^+) D_{\underline{n};n}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda}) \quad (2.154)$$

$$\times \text{Tr}_{V_{1,\infty} \otimes V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty}} P_{\mathcal{L};M}^* Q_{\lambda_*; \underline{\lambda}} B_{\mathcal{L}}[\underline{t}; \underline{\epsilon}] \prod_{j=1}^{L'} \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}'_j}; \beta_j, \pm \infty}} P_{\mathcal{L}';M} Q'_{\underline{\lambda};j} B_{\mathcal{L}'_j; M; \underline{n}}[\underline{t}; \underline{\epsilon}'] \Big|_{\substack{\underline{\epsilon}=0 \\ \lambda_*=0}},$$

where

$$P_{\mathcal{L};M}^* = P_M \otimes P_{\mathcal{L};M}, \quad Q_{\lambda_*; \underline{\lambda}} = \lambda_*^{2\hat{m}} \otimes Q_{\underline{\lambda},*}^{(|\underline{N}|)} \quad (2.155)$$

and

$$Q'_{\underline{\lambda};j} = \begin{cases} (\lambda^{2\hat{m}})^{\otimes |\underline{N}'|} & \text{if } j > 1 \text{ or } \mathcal{L} \text{ is non-empty,} \\ \lambda_0^{2\hat{m}} \otimes (\lambda^{2\hat{m}})^{\otimes N'_j} & \text{if } j = 1 \text{ and } \mathcal{L} \text{ is empty} \end{cases} \quad (2.156)$$

Proof. Equation (2.150) follows from eqs. (2.139) and (2.145). The bounds (2.151) and (2.153) follow from Proposition 2.15 and from eq.(2.148). Equation (2.141) says that its *r.h.s.* represents the expansion of the matrix elements of its *l.h.s.* in powers of h , and according to Proposition 2.11, that expansion leads to the expansion (2.85). Therefore

$$A_{\underline{\beta};n|\underline{n}}(\mathcal{L}, \mathcal{L}'\underline{\lambda}; \underline{t}|M) = F(\underline{n}^+) D_{\underline{n};n}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda}) \quad (2.157)$$

$$\times \text{Tr}_{V_{1,\infty} \otimes V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}'}; \underline{\beta}, \pm \infty}} (P_{\mathcal{L};M}^* \otimes P_{\mathcal{L}';M}) (\lambda_*^{2\hat{m}} \otimes Q_{\underline{\lambda},*}^{(N)}) B_{\mathcal{L} \cup \mathcal{L}'; M; \underline{n}}[\underline{t}; \underline{\epsilon}] \Big|_{\substack{\underline{\epsilon}=0 \\ \lambda_*=0}}$$

According to eq.(2.137), the trace of this equation can be presented as a product of traces, and this leads us to eq.(2.154). $\square$

2.2.5 Calculating the traces

Our next goal is to calculate the traces of eq.(2.154) in the limit of $M \rightarrow \infty$. We start with the trace over $V_{1,\infty} \otimes V_{\mathcal{B}_{\mathcal{L}};\underline{a},\infty}$. We will calculate it with the help of the following

Lemma 2.17 *Let $\hat{O}$ be an operator acting on a space $\mathbb{C}^N$ such that the absolute values of its eigenvalues are less than 1. We extend $\hat{O}$ to a homomorphism O of the algebra $\mathbb{C}[z_1, \dots, z_N]$. Then there exists a limit*

$$\lim_{M \rightarrow \infty} \text{Tr}_{\mathbb{C}[z_1, \dots, z_N]} P_M O = \frac{1}{\det_{\mathbb{C}^N}(1 - \hat{O})}, \quad (2.158)$$

where P_M is a projector onto the polynomials of degree less than M .

Proof. Equation (2.158) is a multi-dimensional generalization of the formula for the sum of a geometric series. $\square$

We identify a Verma module $V_{\alpha,\infty}$ with the algebra $\mathbb{C}[z]$

$$f_m \leftrightarrow z^m. \quad (2.159)$$

Then it is easy to verify that the operators (2.111), (2.112), (2.119) and (2.120) are algebra homomorphisms generated by the following 2×2 matrices

$$\hat{R}^+[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4] = \begin{pmatrix} e^{\epsilon_1 + \epsilon_3}(1 - t_2^{-1} + \epsilon_4) & e^{\epsilon_2} t_1^{-1} \\ e^{\epsilon_1} & 0 \end{pmatrix}, \quad (2.160)$$

$$\hat{R}^-[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4] = \begin{pmatrix} 0 & e^{\epsilon_2} \\ e^{\epsilon_1} t_2 & e^{\epsilon_2 + \epsilon_3} t_1^{-1} t_2 (1 - t_1 + \epsilon_4) \end{pmatrix}, \quad (2.161)$$

$$\hat{R}^{(+)}[\epsilon_1, \epsilon_2] = \begin{pmatrix} 1 & 0 \\ \epsilon_2 & e^{\epsilon_1} \end{pmatrix}, \quad (2.162)$$

$$\hat{R}^{(-)}[\epsilon_1, \epsilon_2] = \begin{pmatrix} 1 & \epsilon_2 e^{\epsilon_1} \\ 0 & e^{\epsilon_1} \end{pmatrix}. \quad (2.163)$$

Therefore, we conclude that the whole matrix $B_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]$ is a homomorphism of the algebra $\mathbb{C}[z_0, \dots, z_{|\underline{N}|}]$ which is generated by a matrix $\hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]$ acting on $\mathbb{C}^{|\underline{N}|+1}$. $\hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]$ is constructed by putting the operators $\hat{R}^{\pm}[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4]$ and $\hat{R}^{(\pm)}[\epsilon_1, \epsilon_2]$ acting on the appropriate

basis vectors of $\mathbb{C}^{|\underline{N}|+1}$, in place of the operators $R^\pm[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4]$ and $R^{(\pm)}[\epsilon_1, \epsilon_2]$ in the expression for $B_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]$.

Since $Q_{\lambda_*; \underline{\lambda}}$ is also an algebra homomorphism generated by the matrix

$$\hat{Q}_{\lambda_*; \underline{\lambda}} = \text{diag}(\lambda_*, \lambda_0, \lambda, \dots, \lambda), \quad (2.164)$$

then we can calculate the first trace of eq.(2.154) in the limit of $M \rightarrow \infty$ with the help of Lemma 2.17.

Proposition 2.18 *There is a vicinity of $\underline{\epsilon} = 0$, $\underline{t} = 1$, $\lambda_*, \underline{\lambda} = 0$ in which there is a uniform convergence*

$$\lim_{M \rightarrow \infty} \text{Tr}_{V_{1, \infty} \otimes V_{\mathcal{B}_{\mathcal{L}}; \underline{\alpha}, \infty}} P_{\mathcal{L}; M}^* Q_{\lambda_*; \underline{\lambda}} B_{\mathcal{L}}[\underline{t}; \underline{\epsilon}] = \frac{1}{\det_{\mathbb{C}^{|\underline{N}|+1}}(I - \hat{Q}_{\lambda_*; \underline{\lambda}} \hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}])}. \quad (2.165)$$

Proof. An inspection the matrix elements of the operators $Q_{\lambda_*; \underline{\lambda}}$ and $B_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]$ indicates that there is a vicinity of $\underline{\epsilon} = 0$, $\underline{t} = 1$, $\lambda_*, \lambda_0, \underline{\lambda} = 0$ in which the absolute values of the eigenvalues of the operator $\hat{Q}_{\lambda_*; \underline{\lambda}} \hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]$ are less than 1. Then eq.(2.165) follows from eq.(2.158). $\square$

We denote

$$Q(\mathcal{L}; \underline{t}; \underline{\lambda}) = \det_{\mathbb{C}^{|\underline{N}|+1}}(I - \hat{Q}_{\lambda_*; \underline{\lambda}} \hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]) \Big|_{\underline{\epsilon}, \lambda_* = 0}, \quad (2.166)$$

because we will show that this determinant is indeed the polynomial mentioned in Proposition 2.7.

Proposition 2.19 *The function $Q(\mathcal{L}; \underline{t}; \underline{\lambda})$ satisfies the properties (2.67) and (2.69).*

Proof. Relation (2.67) follows from the structure of the matrix elements of matrices (2.160)–(2.163) and (2.164). To prove eq.(2.69), note that

$$\hat{R}^\pm[t_1, t_2; \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4] \Big|_{\underline{\epsilon}=0} = \rho_\pm[t_1, t_2], \quad \hat{R}^{(\pm)}[\epsilon_1, \epsilon_2] \Big|_{\epsilon_1, \epsilon_2=0} = I. \quad (2.167)$$

Therefore $\hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}]$ has a block form at $\underline{\epsilon} = 0$

$$\hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}] = \begin{pmatrix} 1 & 0 \\ 0 & B_{\mathcal{L}}[\underline{t}] \end{pmatrix}, \quad (2.168)$$

where $B_{\mathcal{L}}[t]$ is the matrix of Burau representation. Similarly

$$\hat{Q}_{\lambda_*, \Delta} \Big|_{\substack{\lambda_*=0 \\ \lambda=1}} = \begin{pmatrix} 0 & 0 \\ 0 & Q \end{pmatrix}, \quad (2.169)$$

where Q is defined by eq.(1.28). Then eq.(2.69) follows from eq.(1.29). $\square$

Now it remains to calculate the traces

$$\text{Tr}_{V_{\mathcal{B}_{\mathcal{L}'_j; \beta_j, \pm\infty}}} P_{\mathcal{L}'_j; M} Q'_{\Delta; j} B_{\mathcal{L}'_j; M; \underline{n}}[t; \epsilon'] \quad (2.170)$$

of eq. (2.154). Consider the product of matrix elements of operators which constitute $Q'_{\Delta; j} B_{\mathcal{L}'_j; M; \underline{n}}[t; \epsilon']$. Suppose that we start with a vector f_{m_j} of the space V_{β_j} , which is the first space in the second tensor product of (2.42). Since the choice of the numbers $\underline{n}$ fixes the changes in m_j as this vector evolves along the strands of $\mathcal{L}_j$, then it is easy to see that, if $M > \beta_j + 2|\underline{n}|$, then the projectors P_M can be removed from the expression for $B_{\mathcal{L}'_j; M; \underline{n}}[t; \epsilon']$ and the projector $P_{\mathcal{L}'_j; M}$ can be removed from (2.170) without affecting the trace. As a result, the product of the matrix elements has the form $X_j Y_j^{m_j}$ where $X_j(t, \underline{\lambda}, \epsilon')$ and $Y_j(t, \underline{\lambda}, \epsilon')$ are the products of non-negative powers of $e^{\epsilon'}$, $\underline{\lambda}$, $\underline{t}$ and $\underline{t}^{-1}$ which depend on $\underline{n}$ (we do not indicate this dependence explicitly in order to simplify notations). In particular,

$$X_j(t, \underline{\lambda}, \epsilon') \Big|_{\substack{\lambda=1 \\ \lambda_0, \epsilon'=0}} = 1, \quad Y_j(t, \underline{\lambda}, \epsilon') \Big|_{\substack{\lambda=1 \\ \lambda_0, \epsilon'=0}} = \prod_{i=1}^L t_i^{-l_{ij}} \quad \text{if } \underline{n} = 0. \quad (2.171)$$

Calculating the trace over V_{β_j} amounts to taking a sum over m_j , so

$$\lim_{M \rightarrow \infty} \text{Tr}_{V_{\mathcal{B}_{\mathcal{L}'_j; \beta_j, \pm\infty}}} P_{\mathcal{L}'_j; M} Q'_{\Delta; j} B_{\mathcal{L}'_j; M; \underline{n}}[t; \epsilon'] = \sum_{m_j=0}^{\beta_j-1} X_j Y_j^{m_j} = X_j \frac{Y_j^{\beta_j} - 1}{Y_j - 1}. \quad (2.172)$$

Proposition 2.20 *For non-negative integers $l, \underline{l}$ the expression*

$$Z_j(\beta; \underline{t}) = \partial_{\lambda}^l \partial_{\epsilon'}^{\underline{l}} \left(X_j \frac{Y_j^{\beta_j} - 1}{Y_j - 1} \right) \Big|_{\substack{\lambda=1 \\ \lambda_0, \epsilon'=0}} \quad (2.173)$$

has two properties: for fixed β

$$Z_j(\beta; \underline{t}) \in \mathbb{Z}[\underline{t}, \underline{t}^{-1}] \quad (2.174)$$

and its expansion in powers of $\underline{t} - 1$ has the form

$$Z_j(\underline{\beta}; \underline{t}) = \sum_{\underline{m} \geq 0} Z_{j, \underline{m}}(\beta_j) (\underline{t} - 1)^{\underline{m}},$$

$$Z_{j, \underline{m}}(\beta_j) \in \mathbb{Q}[\beta_j], \quad \deg Z_{j, \underline{m}} \leq |\underline{m}| + |\underline{l}| + l + 1. \quad (2.175)$$

Proof. The property (2.174) is easy to see if we reverse eq.(2.172)

$$Z_j(\underline{\beta}; \underline{t}) = \partial_\lambda^l \partial_{\underline{\epsilon}'}^l \left(X_j \sum_{m_j=0}^{\beta_j-1} Y_j^{m_j} \right) \Big|_{\substack{\lambda=1 \\ \lambda_0, \underline{\epsilon}'=0}}. \quad (2.176)$$

In order to prove the property (2.175) we expand the expression (2.172) inside the *r.h.s.* of eq.(2.173) in powers of $Y_j - 1$

$$Z_j(\underline{\beta}; \underline{t}) = \partial_\lambda^l \partial_{\underline{\epsilon}'}^l \left(X_j \sum_{n=0}^{\infty} \binom{\beta_j}{n+1} (Y_j - 1)^n \right) \Big|_{\substack{\lambda=1 \\ \lambda_0, \underline{\epsilon}'=0}}. \quad (2.177)$$

Each of the derivatives $\partial_\lambda, \partial_{\underline{\epsilon}'}$ reduces the power of $Y_j - 1$ at most by one unit. Therefore the bound (2.175) follows from eq.(2.177), if we observe that since $Y_j|_{\substack{\lambda=1 \\ \lambda_0, \underline{\epsilon}'=0}}$ is a product of the powers of $\underline{t}$, then the expansion of $(Y_j - 1)|_{\substack{\lambda=1 \\ \lambda_0, \underline{\epsilon}'=0}}$ in powers of $\underline{t} - 1$ starts at degree 1. $\square$

Now we have all the ingredients which are necessary to finish the proof of Proposition 2.7.

Proof of Proposition 2.7. Combining eqs.(2.172) and (2.165) with eq.(2.154) we conclude that

$$\lim_{M \rightarrow \infty} A_{\underline{\beta}; n | \underline{n}}(\mathcal{L}, \mathcal{L}' \underline{\lambda}; \underline{t} | M) = F(\underline{n}^+) \quad (2.178)$$

$$\times D_{\underline{n}; n}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_\lambda) \frac{\prod_{j=1}^{L'} X_j \frac{Y_j^{\beta_j} - 1}{Y_j - 1}}{\det_{\mathbb{C}^{|\underline{N}|+1}}(I - \hat{Q}_{\lambda*, \underline{\Delta}} \hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}])} \Big|_{\lambda*, \underline{\epsilon}=0}.$$

Since the sum in the *r.h.s.* of eq.(2.109) is finite, then the limit of the *l.h.s.* at $M \rightarrow \infty$ is equal to the sum of limits

$$\lim_{M \rightarrow \infty} A_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t} | M) = \sum_{\substack{\underline{n} \geq 0 \\ |\underline{n}^+| + |\underline{n}^-| \\ |\underline{n}^-| + |\underline{n}''| \leq n}} \left(F(\underline{n}^+) \right. \quad (2.179)$$

$$\left. \times D_{\underline{n}; n}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_\lambda) \frac{\prod_{j=1}^{L'} X_j \frac{Y_j^{\beta_j} - 1}{Y_j - 1}}{\det_{\mathbb{C}^{|\underline{N}|+1}}(I - \hat{Q}_{\lambda*, \underline{\Delta}} \hat{B}_{\mathcal{L}}[\underline{t}; \underline{\epsilon}])} \right) \Big|_{\lambda*, \underline{\epsilon}=0}.$$

The operators $D_{\underline{n};n}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda})$ are combinations of derivatives, their structure is described by Proposition 2.16. The action of the derivatives ∂_{λ} and $\partial_{\underline{\epsilon}}$ on the denominator in the *r.h.s.* of eq.(2.179) will increase its power and bring its derivatives to the numerator. Hence the power of $Q(\mathcal{L}; \underline{t}; \underline{\lambda})$ in the denominator of eq.(2.68) follows from the bound (2.151). The action of the derivatives $\partial_{\underline{\epsilon}'}$, ∂_{λ} on the numerator is described by Proposition 2.20. The order of the derivatives $\partial_{\underline{\epsilon}'}$ at each monomial of $\underline{t} - 1$ in the expansion (2.152) of $D_{\underline{n};n}(\underline{t}; \lambda; \underline{\beta}; \partial_{\underline{\epsilon}}; \partial_{\lambda})$ is restricted by (2.153), hence the bound (2.72).

Thus we proved that the *r.h.s.* of eq.(2.179) has the same form as the *r.h.s.* of eq.(2.68) except that we can only claim that

$$Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}) \in \mathbb{Q}[\underline{t}, \underline{t}^{-1}, \underline{\lambda}]. \quad (2.180)$$

To prove the integrality of the coefficients of $Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda})$ we rewrite eq.(2.68) as

$$Q^{2n+1}(\mathcal{L}; \underline{t}; \underline{\lambda}) \lim_{M \rightarrow \infty} A_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{\lambda}; \underline{t} | M) = Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}). \quad (2.181)$$

Now the integrality of the coefficients of $Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda})$ follows from the integrality of the coefficients of $Q(\mathcal{L}; \underline{t}; \underline{\lambda})$ and from (2.65). Finally, relation (2.70) follows from eqs.(2.150) and (2.171). $\square$

2.3 Removal of parametrization

Now it's time to go back to the original (unparametrized and unregularized) Jones polynomial by setting $\lambda_0 = 0$, $\lambda = 1$ in eq.(2.68). We define a formal power series

$$\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h) = \begin{cases} \frac{1}{h} q^{\phi_{1k}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) + \phi_2(\underline{\beta})} \frac{\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{t})}{\nabla_A(\mathcal{L}; \underline{t})} \sum_{n=0}^{\infty} \frac{Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}) \Big|_{\substack{\lambda_0=0 \\ \lambda=1}}}{\left(\Phi_1^{-1}(\mathcal{B}_{\mathcal{L}}; \underline{t}) (t_1^{1/2} - t_1^{-1/2}) \nabla_A(\mathcal{L}; \underline{t}) \right)^{2n}} h^n, & \text{if } L \geq 1, \\ \frac{1}{h} q^{\phi_{1k}(\emptyset, \mathcal{L}'; \underline{\beta}) + \phi_2(\underline{\beta})} (1 - q^{-\beta_1}) \sum_{n=0}^{\infty} Q_{\underline{\beta};n}(\emptyset; \mathcal{L}'; \underline{\lambda}) \Big|_{\substack{\lambda_0=0 \\ \lambda=1}} h^n & \text{if } L = 0, \end{cases} \quad (2.182)$$

where

$$\phi_2(\underline{\beta}) = \frac{1}{2} \left(\sum_{j=1}^L (l_{jj} - N_j - 1) - \sum_{j=1}^{L'} (l'_{jj} - N'_j - 1)(\beta_j - 1) \right) + 1, \quad (2.183)$$

while $\phi_{\text{lk}}(\mathcal{L}, \mathcal{L}'; \underline{\beta})$ and $\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{t})$ are defined by eqs.(1.60) and (1.61).

We used the same notation for the series (2.182) as in eq.(1.50), because we will ultimately show that both expressions are the same.

Proposition 2.21 *Formal power series (2.182) satisfies three properties of $\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h)$ (eqs. (1.51), (1.52) and (1.53)) listed in Main Theorem, except that if $L \geq 2$, then eq. (1.53) is satisfied only if $i \neq 1$.*

Proof. Equations (1.51), (1.52) follow from eqs.(2.73) and (2.74) respectively if we write the latter equations at $\lambda_0 = 0$, $\lambda = 1$. To prove eq.(1.53) we have to consider two cases.

Suppose that $L \geq 2$ and $i \neq 1$. According to eq.(2.69) and Torres formula (1.24),

$$\begin{aligned} Q(\mathcal{L}; \underline{t}; \underline{\lambda}) \Big|_{\substack{t_i=1 \\ \lambda_0=0 \\ \lambda=1}} &= \Phi_1^{-1}(\mathcal{B}_{\mathcal{L}}; \underline{t}) (t_1^{1/2} - t_1^{-1/2}) \nabla_A(\mathcal{L}; \underline{t}) \Big|_{t_i=1} \\ &= \Phi_1^{-1}(\mathcal{B}_{\mathcal{L}}; \underline{t}) (t_1^{1/2} - t_1^{-1/2}) \Big|_{t_i=1} \left(\prod_{j=1}^{L-1} t_j^{l_{ij}/2} - \prod_{j=1}^{L-1} t_j^{-l_{ij}/2} \right) \nabla_A(\mathcal{L}_{(i)}; \underline{t}_{(i)}) \end{aligned} \quad (2.184)$$

According to our assumptions,

$$\left(\prod_{j=1}^{L-1} t_j^{l_{ij}/2} - \prod_{j=1}^{L-1} t_j^{-l_{ij}/2} \right) \neq 0, \quad \nabla_A(\mathcal{L}_{(i)}; \underline{t}_{(i)}) \neq 0. \quad (2.185)$$

Hence, according to eq.(2.184), $Q(\mathcal{L}; \underline{t}; \underline{\lambda}) \Big|_{\substack{t_i=1 \\ \lambda_0=0 \\ \lambda=1}} \neq 0$ and therefore eq.(1.53) follows from eq.(2.75).

Suppose now that $L = 1$. According to eqs.(2.69), (1.20) and (1.18),

$$Q_0(\mathcal{L}; t_1; \underline{\lambda}) \Big|_{\substack{t_1=1 \\ \lambda_0=0 \\ \lambda=1}} = \Phi_1^{-1}(\mathcal{L}; t_1) \Delta_A(\mathcal{L}; t_1) \Big|_{t_1=1} = 1 \neq 0. \quad (2.186)$$

Therefore we can apply eq.(2.75). Equation (2.64) indicates that

$$\Phi_2(-\alpha_1; \underline{t}_{[1]}; \underline{\lambda}; \underline{\beta}) \Big|_{\lambda_0=0} = 0 \quad \text{if } \alpha_1 > 0, \quad (2.187)$$

so that we can drop the term with $\mu = -1$ from the *l.h.s.* of eq.(2.75) if $\lambda_0 = 0$. Thus we arrive at eq.(1.53) for $L = 1$. $\square$

Proposition 2.22 *There exists a set of polynomials $\tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}) \in \mathbb{Z}[\underline{t}^{1/2}, \underline{t}^{-1/2}, 1/2]$ such that*

$$\tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}) \in \mathbb{Z}[\underline{t}, \underline{t}^{-1}], \quad (2.188)$$

$$\tilde{P}_{\underline{\beta};0}(\mathcal{L}, \mathcal{L}'; \underline{t}) = \prod_{j=1}^{L'} \frac{1 - \left(\prod_{i=1}^L t_i^{l_{ij}}\right)^{-\beta_j}}{1 - \left(\prod_{i=1}^L t_i^{l_{ij}}\right)^{-1}}, \quad (2.189)$$

an expansion of $\tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})$ in powers of $\underline{t} - 1$ has a form

$$\begin{aligned} \tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}) &= \sum_{\underline{m} \geq 0} \tilde{p}_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) (\underline{t} - 1)^{\underline{m}}, \\ \tilde{p}_{\underline{m};n}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) &\in \mathbb{Q}[\underline{\beta}], \quad \deg \tilde{p}_{\underline{m};n} \leq 2(|\underline{m}| + n) + 1. \end{aligned} \quad (2.190)$$

and

$$\begin{aligned} \check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h) & \\ = \begin{cases} q^{\phi_{1k}(\mathcal{L}, \mathcal{L}'; \underline{\beta})} \sum_{n=0}^{\infty} \tilde{P}_{\underline{\beta};n}(\mathcal{L}') h^n & \text{if } L = 0, \\ \frac{1}{h} q^{\phi_{1k}(\mathcal{L}, \mathcal{L}'; \underline{\beta})} \frac{\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; t_1)}{\nabla_A(\mathcal{L}; t_1)} \sum_{n=0}^{\infty} \frac{\tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; t_1)}{\Delta_A^{2n}(\mathcal{L}; t_1)} h^n & \text{if } L = 1, \\ \frac{1}{h} q^{\phi_{1k}(\mathcal{L}, \mathcal{L}'; \underline{\beta})} \frac{\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{t})}{\nabla_A(\mathcal{L}; \underline{t})} \sum_{n=0}^{\infty} \frac{\tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})}{\left((t_1 - 1) \nabla_A(\mathcal{L}; \underline{t})\right)^{2n}} h^n & \text{if } L \geq 2, \end{cases} \end{aligned} \quad (2.191)$$

Proof. The expressions (2.191) can be derived directly from eq.(2.182). If $L = 0$, then we have to expand the factor $q^{\phi_3(\underline{\beta})+1} (1 - q^{-\beta_1})$ in powers of h . If $L \geq 1$, then we expand the factor $q^{\phi_2(\underline{\beta})}$ in powers of h and combine the factors $\Phi_1^{-2n}(\mathcal{B}_{\mathcal{L}}; \underline{t})$ with the polynomials $Q_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}; \underline{\lambda}) \Big|_{\substack{\lambda_0=0 \\ \lambda=1}}$. Equation (2.189) follows from eqs. (2.182), (1.61) and (2.70). The property (2.188) follows from (2.67) and from the fact that $\phi_2(\underline{\beta}) \in \mathbb{Z}$, because it is not hard to prove that

$$l_{jj} - N_j - 1, l'_{jj} - N'_j - 1 \in \mathbb{Z}. \quad (2.192)$$

The bound (2.190) on the degree of the polynomials $\tilde{P}_{\underline{m};n}$ follows easily from the similar bound (2.72). $\square$

3 Uniqueness arguments

3.1 Proof of Main Theorem

The claims of Proposition 2.22 are very similar to those of Main Theorem. In fact we will prove all claims of Main Theorem if we show two things: the formal power series $\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h)$ is an invariant of $\mathcal{L}, \mathcal{L}'$ (that is, its value does not depend on a choice of a braid $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ used to calculate it) and if $L \geq 2$, then polynomials $\tilde{P}_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{t})$ are divisible by $(t_1 - 1)^{2n}$.

Theorem 3.1 (Uniqueness Theorem) *Let $\mathfrak{L}_*$ be a subset of the set of all links, such that the empty link and all 1-component links belong to it, and if $\mathcal{L} \in \mathfrak{L}_*$ and a knot $\mathcal{L}_0$ is algebraically connected to $\mathcal{L}$, then $\mathcal{L}_0 \cup \mathcal{L} \in \mathfrak{L}_*$. Suppose that for any pair of oriented links $\mathcal{L}', \mathcal{L} \in \mathfrak{L}_*$ and for any number l , $1 \leq l \leq L$, there exists a non-empty set $\mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ such that for any $c \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ there exist formal power series*

$$G^{(c)}(\mathcal{L}; \underline{a}) \in \mathbb{Q}[[\underline{a}]], \quad F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}) \in \mathbb{Q}[\underline{\beta}][[\underline{a}]], \quad n \geq 0, \quad (3.1)$$

which satisfy the following properties:

$$G^{(c)}(\mathcal{L}; \underline{a}) \not\equiv 0, \quad (3.2)$$

if $\mathcal{L}_j$ is algebraically connected to $\mathcal{L}_{(j)}$ then

$$G^{(c)}(\mathcal{L}; \underline{a})|_{a_j=0} \not\equiv 0 \quad (3.3)$$

and a formal power series

$$I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = \begin{cases} \sum_{n=0}^{\infty} F_n^{(c)}(\mathcal{L}'; \underline{\beta}) \epsilon^n & \text{if } L = 0, \\ \sum_{n=0}^{\infty} \frac{F_n^{(c)}(\mathcal{L}, \mathcal{L}'; a_1; \underline{\beta})}{(G^{(c)}(\mathcal{L}; a_1))^{2n+1}} \epsilon^n & \text{if } L = 1, \\ \sum_{n=0}^{\infty} \frac{F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})}{(a_l G^{(c)}(\mathcal{L}; \underline{a}))^{2n+1}} \epsilon^n & \text{if } L \geq 2. \end{cases} \quad (3.4)$$

satisfies the following properties

(1) If $\mathcal{L}$ is empty, then $I^{(c)}(\emptyset, \mathcal{L}'; \underline{\beta})$ is an invariant of an oriented link $\mathcal{L}'$, that is, it is the same for all $c \in \mathcal{C}_{\mathcal{L}'}$.

(2) For any j , $1 \leq j \leq L'$ and for any $c \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ there exists $c' \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'_{[1]}; l}$ such that

$$I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) \big|_{\beta_j=1} = I^{(c')}(\mathcal{L}, \mathcal{L}'_{(j)}; \underline{a}; \underline{\beta}_{(j)}; \epsilon). \quad (3.5)$$

(3) Suppose that for a number i , $1 \leq i \leq L$, $\mathcal{L}_{(i)} \in \mathfrak{L}_*$ and $\mathcal{L}_i$ is algebraically connected to $\mathcal{L}_{(i)}$. Then for any $c \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ there exists $c' \in \mathcal{C}_{\mathcal{L}_{[1]}, \mathcal{L}_L \cup \mathcal{L}'; l}$ and for any $c' \in \mathcal{C}_{\mathcal{L}_{[1]}, \mathcal{L}_L \cup \mathcal{L}'; l}$ there exists $c \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ such that

$$\begin{aligned} & I^{(c')}(\mathcal{L}_{(i)}, \mathcal{L}_j \cup \mathcal{L}'; \underline{a}_{(i)}; \beta_0, \underline{\beta}; \epsilon) \\ &= \begin{cases} I^{(c)}(\mathcal{L}, \mathcal{L}'; \epsilon \beta_0; \underline{\beta}; \epsilon) & \text{if } L = 1, \\ \sum_{\mu=\pm 1} \mu e^{i\pi\mu\beta_0 \sum_{\substack{1 \leq j \leq L \\ j \neq i}} l_{ij} a_j} I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) \big|_{a_i=\epsilon\beta_0} & \text{if } L \geq 2 \text{ and } i \neq l. \end{cases} \end{aligned} \quad (3.6)$$

Then

(1) The formal power series $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ is an invariant of a pair consisting of an oriented link $\mathcal{L} \in \mathfrak{L}_*$ and a link $\mathcal{L}'$, that is, for any two numbers l_1, l_2 , $1 \leq l_1, l_2 \leq L$ and for two elements $c_{1,2} \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l_{1,2}}$,

$$I^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = I^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon). \quad (3.7)$$

(2) The invariant $I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ is an odd function of $\underline{\beta}$, that is, for any j , $1 \leq j \leq L'$

$$I(\mathcal{L}, \mathcal{L}'; \underline{a}; \beta_1, \dots, -\beta_j, \dots, \beta_{L'}; \epsilon) = -I(\mathcal{L} \cup \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) \quad (3.8)$$

(we dropped the index (c) , because in view of eq. (3.7), $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ does not depend on it).

(3) If $\mathcal{L}$ is algebraically connected, then the formal power series $F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})$ is divisible by a_l^{2n+1} in $\mathbb{Q}[\underline{\beta}][[\underline{a}]]$.

(4) If the series $G^{(c)}(\mathcal{L}; \underline{a})$ is an invariant of $\mathcal{L}$, then for any link $\mathcal{L} \in \mathfrak{L}_*$ the series $F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})$ is divisible by a_l^{2n+1} in $\mathbb{Q}[\underline{\beta}][[\underline{a}]]$.

Corollary 3.2 *Suppose that in addition to the conditions of Uniqueness Theorem the invariant $I(\emptyset, \mathcal{L}'; \underline{\beta}; \epsilon)$ does not depend on the orientation of $\mathcal{L}'$. Then*

- (1) *The invariant $I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ does not depend on the orientation of $\mathcal{L}'$.*
- (2) *Let $\bar{\mathcal{L}}$ be a link which differs from $\mathcal{L} \in \mathfrak{L}_*$ only by the orientation of its component $\mathcal{L}_j$. Then*

$$I(\bar{\mathcal{L}}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = -I(\mathcal{L} \cup \mathcal{L}'; a_1, \dots, -a_j, \dots, a_L; \underline{\beta}; \epsilon) \quad (3.9)$$

- (3) *If we take an oriented link $\mathcal{L} \in \mathfrak{L}_*$ and reverse the orientations of all of its components, then for the resulting link $\bar{\mathcal{L}}$*

$$I(\bar{\mathcal{L}}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) \quad (3.10)$$

We will prove Uniqueness Theorem and Corollary 3.2 in the next subsection. Here we will show how they help to derive Main Theorem from Proposition 2.22.

Proof of Main Theorem Let $\mathfrak{L}_*$ be a set of all links whose Alexander-Conway function is not identically equal to zero. Let $\mathcal{C}_{\mathcal{L}, \mathcal{L}'; \underline{t}}$ be the set of braids $\mathcal{B}_{\mathcal{L} \cup \mathcal{L}'}$ whose closure is $\mathcal{L} \cup \mathcal{L}'$. Finally, let $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ be a formal power series $h \check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h)$ in which we perform the following substitution

$$\underline{t} = e^{\underline{a}}, \quad h = e^{\epsilon} - 1. \quad (3.11)$$

A simple manipulation with eq.(2.191) brings it to the form (3.4) with

$$G^{(c)}(\mathcal{L}; \underline{a}) = \begin{cases} \Delta_A(\mathcal{L}; e^{a_1}) & \text{if } L = 1, \\ \nabla_A(\mathcal{L}; e^{\underline{a}}) & \text{if } L \geq 2. \end{cases} \quad (3.12)$$

Now a simple manipulation brings $\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h)$ of eq. (2.191) to the form (3.4). Therefore Proposition 2.22 implies that all conditions of Uniqueness Theorem hold. In its turn, Uniqueness Theorem says that for $L \geq 2$, $\tilde{P}_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; e^{\underline{a}})$ has to be divisible by a_1^{2n} . In view of (2.188) this means that

$$\frac{\tilde{P}_{\underline{\beta}; n}(\mathcal{L}, \mathcal{L}'; \underline{t})}{(t_1 - 1)^{2n}} \in \mathbb{Z}[\underline{t}, \underline{t}^{-1}]. \quad (3.13)$$

Thus we can set

$$P'_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}) = \begin{cases} \tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t}) & \text{if } L = 0, 1, \\ \frac{\tilde{P}_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})}{(t_1 - 1)^{2n}} & \text{if } L \geq 2, \end{cases} \quad (3.14)$$

and eq.(2.191) will turn into eq.(1.59) with the relation (3.12) guaranteeing (1.62).

Equation (1.50) follows from eq.(1.59) if we combine $\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{t})$ with the polynomials $P'_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})$ and expand the factor $q^{\phi_{\text{lk}}(\mathcal{L}, \mathcal{L}'; \underline{\beta})}$ in powers of h . Equation (1.54) follows from eq.(2.189). Relation (1.48) follows from (1.62) because

$$\Phi_{\underline{\beta}}(\mathcal{L}, \mathcal{L}'; \underline{t}) \in \mathbb{Z}[\underline{t}^{1/2}, \underline{t}^{-1/2}], \quad \phi_{\text{lk}}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \in \frac{1}{2} \mathbb{Z}. \quad (3.15)$$

The property (1.51) of $\check{J}_{\underline{\beta}}^{(r)}(\emptyset, \mathcal{L}'; h)$ together with the independence of the colored Jones polynomial $J_{\underline{\beta}}(\mathcal{L}'; q)$ of the orientation of the link $\mathcal{L}'$ implies that the condition of Corollary 3.2 is satisfied. Therefore Uniqueness Theorem and Corollary (3.2) together say that the formal power series $\check{J}_{\underline{\beta}}^{(r)}(\mathcal{L}, \mathcal{L}'; \underline{t}; h)$ (and hence the polynomials $P_{\underline{\beta};n}(\mathcal{L}, \mathcal{L}'; \underline{t})$) is an invariant of an oriented link $\mathcal{L}$ and a link $\mathcal{L}'$.

Relations (1.55), (1.56) follow from eqs.(3.9), (3.10) and from the properties (1.22), (1.23) of the Alexander-Conway function. Relation (1.57) follows from eq.(3.8). $\square$

3.2 Proof of Uniqueness Theorem

3.2.1 Algebraically connected links

First of all, we will prove an obvious lemma.

Lemma 3.3 *The set $\mathfrak{L}_*$ of Uniqueness Theorem contains all algebraically connected links.*

Proof. We use induction in the number L of link components. Since $\mathfrak{L}_*$ contains the empty link, then lemma holds for $L = 0$. Suppose that all the algebraically connected links with $L - 1$ components belong to $\mathfrak{L}_*$. Let $\mathcal{L}$ be an L -component algebraically connected link. Then $\mathcal{L}_{[1]} \in \mathfrak{L}_*$. Since, by definition, $\mathcal{L}_L$ is algebraically connected to $\mathcal{L}_{[1]}$, then, according to the property of $\mathfrak{L}_*$, $\mathcal{L} \in \mathfrak{L}_*$. $\square$

Proposition 3.4 *Claims 1 and 3 of Uniqueness Theorem hold for algebraically connected links.*

We need two lemmas in order to prove this proposition.

Lemma 3.5 *If the series $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ is unique for an L -component algebraically connected link $\mathcal{L}$ with $L \geq 2$, then $F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})$ is divisible by a_l^{2n+1} in $\mathbb{Q}[\underline{\beta}][[\underline{a}]]$.*

Proof. For a number l , $1 \leq l \leq L$ and an element $c \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ choose another number $l' \neq l$, $1 \leq l' \leq L$ and an element $c' \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l'}$. Uniqueness means that for any $n \geq 0$

$$\frac{F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})}{\left(a_l G^{(c)}(\mathcal{L}; \underline{a})\right)^{2n+1}} = \frac{F_n^{(c')}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})}{\left(a_{l'} G^{(c')}(\mathcal{L}; \underline{a})\right)^{2n+1}}, \quad (3.16)$$

or, equivalently,

$$F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}) \left(a_{l'} G^{(c')}(\mathcal{L}; \underline{a})\right)^{2n+1} = F_n^{(c')}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}) \left(a_l G^{(c)}(\mathcal{L}; \underline{a})\right)^{2n+1}. \quad (3.17)$$

The *r.h.s.* of this equation is divisible by a_l^{2n+1} , while $G^{(c')}(\mathcal{L}; \underline{a})$ is not divisible by a_l , because, since $\mathcal{L}$ is algebraically connected, then $\mathcal{L}_l$ is algebraically connected to $\mathcal{L} \setminus \mathcal{L}_l$ and as a result, according to eq.(3.3),

$$G^{(c')}(\mathcal{L}; \underline{a}) \Big|_{a_l} \neq 0. \quad (3.18)$$

Therefore $F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})$ has to be divisible by a_l^{2n+1} . $\square$

Lemma 3.6 *Let $L \geq 2$. Assume that the series $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ is unique for all algebraically connected $L - 1$ -component links. Suppose that the components of an L -component link $\mathcal{L}$ are indexed in such a way that the link $\mathcal{L}_{[1]}$ is also algebraically connected. Then for any two numbers $l_1, l_2 \neq L$ and for any two elements $c_1 \in \mathcal{C}_{\mathcal{L} \cup \mathcal{L}'; l_1}$, $c_2 \in \mathcal{C}_{\mathcal{L} \cup \mathcal{L}'; l_2}$*

$$I^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = I^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon). \quad (3.19)$$

Proof. Since $\mathcal{L}$ is algebraically connected, then $\mathcal{L}_L$ is algebraically connected to $\mathcal{L}_{[1]}$ and then, according to eq.(3.3),

$$G^{(c_i)}(\mathcal{L}; \underline{a})|_{a_L=0} \neq 0, \quad i = 1, 2. \quad (3.20)$$

Therefore we can expand $I^{(c_i)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ in powers of a_L

$$I^{(c_i)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = \sum_{m,n=0}^{\infty} I_{m,n}^{(c_i)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}) a_L^m \epsilon^n, \quad i = 1, 2, \quad (3.21)$$

$$I_{m,n}^{(c_i)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}) = \frac{H_{m,n}^{(c_i)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta})}{a_{l_i}^{2n+1} \left(G^{(c_i)}(\mathcal{L}; \underline{a}) \right)^{2n+m+1}}, \quad (3.22)$$

where $H_{m,n}^{(c_i)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}) \in \mathbb{Q}[\underline{\beta}][\underline{a}_{[1]}]$. Equation (3.19) would follow from the relations

$$I_{m,n}^{(c_1, c_2)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}) = 0, \quad m, n \geq 0, \quad (3.23)$$

where

$$I_{m,n}^{(c_1, c_2)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}) = I_{m,n}^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}) - I_{m,n}^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}). \quad (3.24)$$

We are going to prove relations (3.23). Since, according to our assumptions, $\mathcal{L}_{[1]}$ is algebraically connected and the series $I^{(c)}(\mathcal{L}_{[1]}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta})$ is unique, then, in view of eq.(3.6),

$$\sum_{\mu_L = \pm 1} \mu_L e^{i\pi \mu_L \alpha_L \sum_{j=1}^{L-1} l_{jL} a_j} \left(I^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \mu_L \alpha_L; \underline{\beta}; \epsilon) - I^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \mu_L \alpha_L; \underline{\beta}; \epsilon) \right) = 0. \quad (3.25)$$

Rewriting the series $I^{(c_i)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \mu_L \alpha_L; \underline{\beta}; \epsilon)$ with the help of eq.(3.21), we find that

$$\sum_{n=0}^{\infty} \epsilon^n \left[\sum_{\mu_L = \pm 1} \mu_L e^{i\pi \mu_L \alpha_L \sum_{j=1}^{L-1} l_{jL} a_j} \left(\sum_{m=0}^n I_{m, n-m}^{(c_1, c_2)}(\mathcal{L}, \mathcal{L}'; \underline{a}_{[1]}; \underline{\beta}) \mu_L^m \alpha_L^m \right) \right] = 0. \quad (3.26)$$

As a function of α_L , each of the two terms in the sum over μ_L is a product of an exponential and a polynomial. Since, according to our assumptions,

$$\mu_L \alpha_L \sum_{j=1}^{L-1} l_{jL} a_j \neq 0, \quad (3.27)$$

then the exponents in both terms are different and therefore each term in the sum over μ_L has to be equal to zero. Hence we come to eq.(3.23) and to eq.(3.19). $\square$

Proof of Proposition 3.4. The proof is by induction in the number of link components L .

If $L = 0$, then the claim of Uniqueness Theorem is equivalent to Property 1 of the series $I^{(c)}(\mathcal{L}'; \underline{\beta})$.

If $L = 1$, then the uniqueness of $I^{(c)}(\mathcal{L}_1, \mathcal{L}'; a_1; \underline{\beta})$ follows from eq. (3.6) and from the uniqueness of $I^{(c')}(\mathcal{L}'; \epsilon a_1, \underline{\beta})$.

If $L = 2$, then we have to consider two cases. For $l = 1, 2$ and for $c_{1,2} \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ the uniqueness relation (3.6) follows from Lemma 3.6. The case of $l_1 = 1, l_2 = 2$ requires a special consideration. Let us expand

$$\frac{F_n^{(c_j)}(\mathcal{L}, \mathcal{L}'; a_1, a_2; \underline{\beta})}{(G^{(c_j)}(\mathcal{L}; a_1, a_2))^{2n+1}} = \sum_{m_1, m_2=0}^{\infty} H_{m_1, m_2, n}^{(c_j)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) a_1^{m_1} a_2^{m_2}. \quad (3.28)$$

Then the uniqueness of the series $I^{(c)}(\mathcal{L}, \mathcal{L}'; a_1, a_2; \underline{\beta})$ would follow from the relation

$$H_{2n+m_1+1, m_2, n}^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) = H_{m_1, 2n+m_2+1, n}^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \quad \text{for all } m_1, m_2 \text{ and } n. \quad (3.29)$$

Let us apply the property (3.6) to $I^{(c_j)}(\mathcal{L}, \mathcal{L}'; a_1, a_2; \underline{\beta})$ twice first reducing it to $I^{(c'_j)}(\mathcal{L}_j, \mathcal{L}_k \cup \mathcal{L}'; a_j; \epsilon \alpha_k, \underline{\beta})$, where $1 \leq k \neq j \leq 2$, and then reducing it further to $I^{(c''_j)}(\emptyset; \mathcal{L} \cup \mathcal{L}'; \epsilon \alpha_1, \epsilon \alpha_2, \underline{\beta})$. We introduce a notation

$$\begin{aligned} \sum_{m_1, m_2, n} X_{m_1, m_2, n}^{(c_1, c_2)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \alpha_1^{m_1} \alpha_2^{m_2} \epsilon^n &= \sum_{\mu=\pm 1} \mu e^{i\pi \epsilon l_{12} \alpha_1 \alpha_2} \\ &\times \sum_{m_1, m_2, n=0}^{\infty} \left(H_{m_1, m_2, n}^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \mu^{m_2} \alpha_1^{m_1-2n-1} \alpha_2^{m_2} \right. \\ &\quad \left. + H_{m_1, m_2, n}^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \mu^{m_1} \alpha_1^{m_1} \alpha_2^{m_2-2n-1} \right) \epsilon^{m_1+m_2-n-1} \end{aligned} \quad (3.30)$$

(note that only a finite number of coefficients $H_{m_1, m_2, n}^{(c_j)}(\mathcal{L}, \mathcal{L}'; \underline{\beta})$ contributes to any given coefficient X). The uniqueness of $I^{(c'')}(\emptyset; \mathcal{L} \cup \mathcal{L}'; \epsilon \alpha_1, \epsilon \alpha_2, \underline{\beta})$ implies the following relation

$$X_{m_1, m_2, n}^{(c_1, c_2)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) = 0 \quad \text{for all } m_1, m_2, n. \quad (3.31)$$

It is easy to see from eq. (3.30) that for any two integers m_1, m_2

$$\begin{aligned} \sum_n X_{m_1-n, m_2-n, m_1+m_2-n}^{(c_1, c_2)} \alpha_1^{m_1-n} \alpha_2^{m_2-n} \epsilon^{m_1+m_2-n} &= \sum_{\mu=\pm 1} \mu e^{i\pi \epsilon l_{12} \alpha_1 \alpha_2} \\ &\times \sum_n \left(H_{m_1+n-1, m_2-n, n}^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) - H_{m_1-n, m_2+n-1, n}^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) \right) \alpha_1^{m_1-n} \alpha_2^{m_2-n} \epsilon^{m_1+m_2-n}. \end{aligned} \quad (3.32)$$

Therefore, the relation (3.31) implies that the *r.h.s.* of eq.(3.32) has to be identically equal to zero. Since the indices of the coefficients H have to be non-negative, then the sum over n in eq.(3.32) contains only finitely many terms. Therefore since $l_{1,2} \neq 0$, then the *r.h.s.* of eq.(3.32) can be identically equal to zero only if

$$H_{m_1+n+1, m_2-n, n}^{(c_1)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) - H_{m_1-n, m_2+n+1, n}^{(c_2)}(\mathcal{L}, \mathcal{L}'; \underline{\beta}) = 0 \quad \text{for any } m_1, m_2 \text{ and } n. \quad (3.33)$$

This condition is obviously equivalent to (3.29), hence we proved the uniqueness of the series $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ for 2-component links.

Suppose now that $L > 2$ and that the series $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ is unique for all algebraically connected links with $L - 1$ components. Consider an L -component algebraically connected link $\mathcal{L}$. Let us construct a special graph $\Gamma(\mathcal{L})$ with L vertices identified with link components. Two vertices corresponding to link components $\mathcal{L}_{j_1}$ and $\mathcal{L}_{j_2}$ are connected by an edge if there exists a third component $\mathcal{L}_k$ such that $k \neq j_1, j_2$ and the sublink $\mathcal{L} \setminus \mathcal{L}_k$ is algebraically connected.

Lemma 3.7 *For an algebraically connected link $\mathcal{L}$, the graph $\Gamma(\mathcal{L})$ is connected.*

Proof. We leave the proof to the reader. □

According to Lemma 3.6, if two vertices j_1 and j_2 of the graph are connected, then for any two elements $c_1 \in \mathcal{C}_{\mathcal{L} \cup \mathcal{L}'; j_1}$, $c_2 \in \mathcal{C}_{\mathcal{L} \cup \mathcal{L}'; j_2}$ the corresponding series $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ are equal. Therefore, the uniqueness of this series for $\mathcal{L}$ follows from Lemma 3.7. This completes the induction argument for the proof of uniqueness of $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ for algebraically connected links. The divisibility of $F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta})$ by a_l^{2n+1} follows from Lemma 3.5. □

3.2.2 General links

Now we can prove the claims of Uniqueness Theorem for any link $\mathcal{L} \in \mathfrak{L}_*$.

Proof Claim 1 of Uniqueness Theorem. Let $\mathcal{L} \in \mathfrak{L}_*$ be an L -component link. Let us add an extra 0-th component $\mathcal{L}_0$ such that the link $\mathcal{L}_0 \cup \mathcal{L}$ is algebraically connected. According to

eq.(3.6), for any integer l , $1 \leq l \leq L$ and for any $c \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ there exists $c' \in \mathcal{C}_{\mathcal{L}_0, \mathcal{L}'; l}$ such that

$$I^{(c)}(\mathcal{L}, \mathcal{L}_0 \cup \mathcal{L}'; \underline{a}; \alpha_0, \underline{\beta}; \epsilon) = \sum_{\mu_0 = \pm 1} \mu_0 e^{i\pi \mu_0 \alpha_0 \sum_{j=1}^{L-1} l_{jL} a_j} I^{(c')}(\mathcal{L}_0 \cup \mathcal{L}, \mathcal{L}'; \mu_0 \alpha_0, \underline{a}; \underline{\beta}; \epsilon). \quad (3.34)$$

Equation (3.5) allows us to remove the extra component $\mathcal{L}_0$ from the *l.h.s.* of eq.(3.34)

$$I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = \sum_{\mu_0 = \pm 1} \mu_0 e^{i\pi \mu_0 \alpha_0 \sum_{j=1}^{L-1} l_{jL} a_j} I^{(c')}(\mathcal{L}_0 \cup \mathcal{L}, \mathcal{L}'; \mu_0 \alpha_0, \underline{a}; \underline{\beta}; \epsilon) \Big|_{\alpha_0=1}. \quad (3.35)$$

The link $\mathcal{L}_0 \cup \mathcal{L}$ is algebraically connected. Therefore, according to Proposition 3.4, the *r.h.s.* of eq.(3.35) does not depend on c' , hence the *l.h.s.* does not depend on c . This proves the uniqueness of $I^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$. $\square$

Proof of Claim 2 of Uniqueness Theorem. Consider an oriented L -component link $\mathcal{L} \in \mathfrak{L}_*$ and an oriented L' -component link $\mathcal{L}'$. Choose a number k , $1 \leq k \leq L'$. Let us add an extra component $\mathcal{L}_0$ to $\mathcal{L}$ in such a way that $l''_{0k} = 1$ and, in addition, $l_{0j} = 1$ for all j , $1 \leq j \leq L$. It is easy to see that both $\mathcal{L}_0 \cup \mathcal{L}$ and $\mathcal{L}'_k \cup \mathcal{L}_0 \cup \mathcal{L}$ are algebraically connected and therefore $\mathcal{L}_0 \cup \mathcal{L}, \mathcal{L}'_k \cup \mathcal{L}_0 \cup \mathcal{L} \in \mathfrak{L}_*$.

According to eq.(3.6),

$$\begin{aligned} & I(\mathcal{L}_0 \cup \mathcal{L}, \mathcal{L}'; a_0, \underline{a}; \underline{\beta}; \epsilon) \\ &= \sum_{\mu = \pm 1} \mu e^{i\pi \mu \beta_k (a_0 + \sum_{j=1}^L l''_{jk} a_j)} I(\mathcal{L}'_k \cup \mathcal{L}_0 \cup \mathcal{L}, \mathcal{L}'_{(k)}; \mu \epsilon \beta_k, a_0, \underline{a}; \underline{\beta}_{(k)}; \epsilon). \end{aligned} \quad (3.36)$$

On the other hand, a combination of eqs.(3.5) and (3.6) implies that

$$I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = \sum_{\mu = \pm 1} \mu e^{i\pi \mu \alpha_0 \sum_{j=1}^L l_{0j} a_j} I(\mathcal{L}_0 \cup \mathcal{L}, \mathcal{L}'; \mu \epsilon \alpha_0, \underline{a}; \underline{\beta}; \epsilon) \Big|_{\alpha_0=1}. \quad (3.37)$$

Equation (3.36) indicates that $I(\mathcal{L}_0 \cup \mathcal{L}, \mathcal{L}'; a_0, \underline{a}; \underline{\beta}; \epsilon)$ is an odd function of β_k . Then it follows from eq.(3.37), that $I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ is also an odd function of β_k . $\square$

Proof of Claim 4 of Uniqueness Theorem. For a number l , $1 \leq l \leq L$ and an element $c \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l}$ choose another number $l' \neq l$, $1 \leq l' \leq L$ and an element $c' \in \mathcal{C}_{\mathcal{L}, \mathcal{L}'; l'}$. Uniqueness means that for any $n \geq 0$ eq.(3.17) holds. If $G^{(c)}(\mathcal{L}; \underline{a})$ is unique (that is, if it does not depend on l and c), then we can cancel it from both sides of that equation, so that

$$F_n^{(c)}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}) a_{l'}^{2n+1} = F_n^{(c')}(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}) a_l^{2n+1}. \quad (3.38)$$

Divisibility follows from this equation. $\square$

3.3 Proof of Corollary 3.2

Proof of Claims 1 and 2 of Corollary 3.2. Let $\mathcal{L} \in \mathfrak{L}_*$ be an L -component oriented link and let $\mathcal{L}'$ be an oriented L' -component link. Consider a group $G_{\mathcal{L}, \mathcal{L}'}$ which is generated by L elements g_j , $1 \leq j \leq L$ each of which acts on $\mathcal{L}$ by changing the orientation of the corresponding link component $\mathcal{L}_j$ and on $\underline{a}$ by changing the sign of a_j , and by L' elements g'_j , $1 \leq j \leq L'$ each of which acts on $\mathcal{L}'$ by changing the orientation of the corresponding link component $\mathcal{L}'_j$. For $g \in G_{\mathcal{L}, \mathcal{L}'}$ let $|g|$ denote the number of components of $\mathcal{L}$ whose orientation it changes.

Consider a new invariant

$$I_S(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = 2^{-(L+L')} \sum_{g \in G_{\mathcal{L}, \mathcal{L}'}} (-1)^{|g|} I(g(\mathcal{L}, \mathcal{L}'); g(\underline{a}), \underline{\beta}; \epsilon). \quad (3.39)$$

It is not hard to see that $I_S(\mathcal{L} \cup \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ satisfies the conditions of Uniqueness Theorem (note that condition (3.6) at $L = 1$ follows from Claim 2 of Uniqueness Theorem applied to $I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$). Since we assume that $I(\emptyset, \mathcal{L}'; \underline{\beta}; \epsilon)$ does not depend on the orientation of $\mathcal{L}'$, then

$$I_S(\emptyset; \mathcal{L}'; \underline{\beta}; \epsilon) = I(\emptyset, \mathcal{L}'; \underline{\beta}; \epsilon). \quad (3.40)$$

Therefore Uniqueness Theorem says that

$$I_S(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon). \quad (3.41)$$

Since the *r.h.s.* of eq.(3.39) satisfies Claims 1 and 2 of Corollary 3.2, then so does the *r.h.s.* of eq.(3.41). $\square$

Proof of Claim 3 of Corollary 3.2. The proof is similar to the previous one. For an oriented link $\mathcal{L}$ let $\bar{\mathcal{L}}$ denote the same link with reversed orientation of all its compomenents. This time we define the new invariant $I_S(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ as

$$I_S(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) = \frac{1}{2} \left(I(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) + I(\bar{\mathcal{L}}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon) \right). \quad (3.42)$$

It follows easily from Claim 1 of Corollary 3.2 that $I_S(\mathcal{L}, \mathcal{L}'; \underline{a}; \underline{\beta}; \epsilon)$ satisfies conditions of Uniqueness Theorem and eq.(3.40). Therefore, according to Uniqueness Theorem, eq.(3.41)

holds, and since the *r.h.s.* of eq.(3.42) satisfies Claim 3 of Corollary 3.2, then so does the *r.h.s.* of eq.(3.41). $\square$

Appendix

A Non-abelian connections in link complements

In this appendix we will prove Theorem 1.13. First, we have to establish two simple lemmas.

Lemma A.1 *Let $\mathcal{L}$ be an algebraically connected L -component link with admissible indexing of its components. Then there exist positive numbers $\underline{c} = (c_1, \dots, c_L)$ and positive integers $\underline{n} = (n_1, \dots, n_{L-1})$ such that $\nabla_A(\mathcal{L}; e^{2\pi i \underline{a}}) \neq 0$ for any $\underline{a} \in \mathcal{A}_{\underline{c}, \underline{n}}$.*

Proof. We will prove the lemma by induction in L . If $L = 1$, then the claim of the lemma follows from eqs.(1.18) and (1.20). Now suppose that $L \geq 2$ and that the lemma holds for $(L - 1)$ -component links. Let us expand $\nabla_A(\mathcal{L}; \underline{t})$ in powers of $t_L - 1$. The first term in this expansion is described by the Torres formula (1.24), so

$$\nabla_A(\mathcal{L}; \underline{t}) = \left(\prod_{j=1}^{L-1} t_j^{l_{jL}/2} - \prod_{j=1}^{L-1} t_j^{-l_{jL}/2} \right) \nabla_A(\mathcal{L}_{(L)}; \underline{t}_{(L)}) + (t_L - 1)F(\mathcal{L}; \underline{t}), \quad (\text{A.1})$$

$$F(\mathcal{L}; \underline{t}) \in \mathbb{Z}[\underline{t}^{1/2}, \underline{t}^{-1/2}].$$

Now we substitute $\underline{t} = e^{2\pi i \underline{a}}$ in eq.(A.1). It is easy to see that there exists an area of the form (1.106) in which

$$\left| \left(\prod_{j=1}^{L-1} e^{i\pi l_{jL} a_j} - \prod_{j=1}^{L-1} e^{-i\pi l_{jL} a_j} \right) \nabla_A(\mathcal{L}_{(L)}; e^{2\pi i \underline{a}}) \right| > |(e^{2\pi i a_L} - 1)F(\mathcal{L}; e^{2\pi i \underline{a}})|. \quad (\text{A.2})$$

Since $\mathcal{L}_L$ is algebraically connected to $\mathcal{L}_{(L)}$, then there exists another area of the form (1.106) in which

$$\prod_{j=1}^{L-1} e^{i\pi l_{jL} a_j} - \prod_{j=1}^{L-1} e^{-i\pi l_{jL} a_j} \neq 0. \quad (\text{A.3})$$

Finally, according to the assumption of induction, there exists an area of the form (1.106) in which

$$\nabla_A(\mathcal{L}_{(L)}; e^{2\pi i \underline{a}_{(L)}}) \neq 0. \quad (\text{A.4})$$

According to eq.(A.1), the intersection of these three areas is the area of the form (1.106) in which $\nabla_A(\mathcal{L}; e^{2\pi i \underline{a}}) \neq 0$. $\square$

The second lemma says that the Alexander-Conway function measures an obstruction to an irreducible deformation of a $U(1)$ -reducible connection. The proof is based on the observation that the Alexander matrix gives a system of linear equations governing this deformation in the linear approximation.

Let G be a simple Lie group, $\mathfrak{g}$ be its Lie algebra. Let H be a maximal torus of G and let $\mathfrak{h}$ be the corresponding Cartan subalgebra. We denote by $\Delta_+ \subset \mathfrak{h}^*$ a set of positive roots of $\mathfrak{g}$.

Let $\mathcal{L}$ be an L -component link in S^3 . We denote by π_1 the group of $\mathcal{L}$ which is the fundamental group of its complement: $\pi_1 = \pi_1(S^3 \setminus \mathcal{L})$. A flat connection A of the trivial G -bundle over $S^3 \setminus \mathcal{L}$ is determined by the corresponding homomorphism

$$f_A : \pi_1 \rightarrow G \quad (\text{A.5})$$

up to an overall conjugation. The homomorphism (A.5) is, of course, generated by the holonomies of the flat connection along the elements of π_1 . We call a flat connection H -reducible, if there exists a maximal torus $H \subset G$ such that the homomorphism (B.1) maps π_1 into H :

$$f_A : \pi_1 \rightarrow H \subset G \quad (\text{A.6})$$

Since H is abelian, then such a homomorphism is determined by the images of the meridians of the link components

$$f_A : \text{mer}(\mathcal{L}_j) \mapsto \exp(x_j), \quad \underline{x} \in \mathfrak{h} \quad (\text{A.7})$$

up to the action of the Weyl group.

Lemma A.2 *Consider a flat H -reducible connection in the G -bundle over the link complement $S^3 \setminus \mathcal{L}$, determined by eqs. (A.6) and (A.7). Suppose that for any positive root $\lambda \in \Delta_+$:*

(1) *if $L = 1$, then*

$$\Delta_A(\mathcal{L}; e^{(\lambda, \underline{x})}) \neq 0; \quad (\text{A.8})$$

(2) *if $L \geq 2$, then*

$$\nabla_A(\mathcal{L}; e^{(\lambda, \underline{x})}) \neq 0, \quad (\text{A.9})$$

and for at least one value of j

$$(\lambda, x_j) \neq 0. \quad (\text{A.10})$$

Here $(\cdot, \cdot)$ denotes a natural pairing between the elements of $\mathfrak{h}$ and its dual $\mathfrak{h}^*$. Then any infinitesimal deformation of this connection should also be H -reducible.

Proof. First of all, we recall the Wirtinger presentation of π_1 (see, *e.g.* Chapter 3 of [2]). The link $\mathcal{L}$ is projected onto a plane, so that each of its components is split by overcrossings into arcs. We denote the number of arcs for $\mathcal{L}_j$ as N_j . Then π_1 is generated by $\underline{N} = \sum_{j=1}^L N_j$ generators s_i which are meridians of the arcs. We introduce a function $l(i)$ such that s_i is the meridian of an arc which is a part of $\mathcal{L}_{l(i)}$. We can always choose a projection of $\mathcal{L}$ in such a way that $\underline{N}$ equals the number of crossings in that projection. Then the meridians s_i , as elements of π_1 , are related by $\underline{N}$ equations

$$r_i = 1, \quad 1 \leq i \leq \underline{N}, \quad (\text{A.11})$$

where the relators r_i come from the crossings in the projection of $\mathcal{L}$ and have a form

$$r_i = s_i s_k^{-1} s_m s_k \quad \text{or} \quad r_i = s_i s_k s_m^{-1} s_k^{-1}, \quad l(i) = l(m), \quad (\text{A.12})$$

depending on the type of the i -th crossing.

A homomorphism f_A describing a flat H -reducible connection maps the elements s_i as

$$f_A(s_i) = \exp(x_{l(i)}). \quad (\text{A.13})$$

Consider an infinitesimal deformation of this map

$$f_A(s_i) = \exp(\epsilon_i) \exp(x_{l(i)}), \quad (\text{A.14})$$

where ϵ_i are infinitesimal elements of $\mathfrak{g}$. The images of the elements s_i should satisfy the relations (A.12). It is easy to see that to the first order in ϵ this means that for any relator r_i ,

$$f_A \left(\frac{\partial r_i}{\partial s_i} \right) (\epsilon_i) + f_A \left(\frac{\partial r_i}{\partial s_k} \right) (\epsilon_k) + f_A \left(\frac{\partial r_i}{\partial s_m} \right) (\epsilon_m) = 0. \quad (\text{A.15})$$

Here $(\partial r / \partial s)$ denote Fox derivatives (see *e.g.* [2], Chapter 9,B) and for a group element $g \in G$ and Lie algebra element $\epsilon \in \mathfrak{g}$, $g(\epsilon)$ denotes the adjoint action of g on ϵ .

Let us decompose ϵ_i into sums of Cartan algebra elements and root space elements. Since the flat connection A is H -reducible, then equations (A.15) do not mix different spaces. Therefore, since we investigate the existence of irreducible deformations, then we may assume that all ϵ_i belong to the same root space $V_\lambda \subset \mathfrak{g}$ corresponding to a root λ :

$$\epsilon_i = b_i \epsilon, \quad b_i \in \mathbb{R}, \quad \epsilon \in V_\lambda. \quad (\text{A.16})$$

Let $f_{A,\lambda} : \pi_1 \rightarrow \mathbb{C}$ be a homomorphism such that

$$f_{A,\lambda}(s_i) = e^{(\lambda, x_{l(i)})}, \quad 1 \leq i \leq \underline{N} \quad (\text{A.17})$$

($f_{A,\lambda}$ is well-defined, because $f_{A,\lambda}(r_i) = 1$). Then the system of equations (A.15) for elements (A.16) can be rewritten as

$$f_{A,\lambda} \left(\frac{\partial r_i}{\partial s_i} \right) b_i + f_{A,\lambda} \left(\frac{\partial r_i}{\partial s_k} \right) b_k + f_{A,\lambda} \left(\frac{\partial r_i}{\partial s_m} \right) b_m = 0, \quad 1 \leq i \leq \underline{N}. \quad (\text{A.18})$$

If $L = 1$, then according to Lemma 9.12 of [2], the matrix of equations (A.18) is degenerate and has a solution of the form

$$b_i = b, \quad 1 \leq i \leq \underline{N}, \quad b \in \mathbb{R}. \quad (\text{A.19})$$

This is the only solution, because, according to Theorem 9.10 of [2], the determinant of any $(\underline{N} - 1) \times (\underline{N} - 1)$ minor of the matrix of (A.18) is proportional to $\Delta_A(\mathcal{L}; e^{(\lambda, \underline{x})})$, so it is not equal to zero in view of condition (1.27).

If $L \geq 2$, then according to the proof of Proposition 9.16 of [2], the matrix of equations (A.18) is degenerate and has a solution of the form

$$b_i = (e^{(\lambda, x_{l(i)})} - 1) b, \quad 1 \leq i \leq \underline{N}, \quad b \in \mathbb{R}. \quad (\text{A.20})$$

This is the only solution, because the determinant of an $(\underline{N} - 1) \times (\underline{N} - 1)$ minor of the matrix of (A.18) obtained by removing the i -th column and the j -th row, is proportional to $(e^{(\lambda, x_{l(i)})} - 1) \nabla_A(\mathcal{L}; e^{(\lambda, \underline{x})})$, the coefficients being the powers of $e^{(\lambda, \underline{x})}$, so at least some minors are not equal to zero in view of conditions (A.9) and (A.10).

The deformation (A.14) with ϵ_i given by either eq. (A.19) or by eqs. (A.16) and (A.20) is in fact H -reducible, because it is easy to see that $f_A(s_i)$ commute with each other in the first order in ϵ and therefore can be conjugated back into H at this order. $\square$

Let us introduce the following notation: if for two Lie group elements $g_1, g_2 \in G$ there exists $h \in G$ such that $g_2 = hg_1h^{-1}$, then we denote this as $g_1 \sim g_2$.

Now we can prove the following

Theorem A.3 *Let $\mathcal{L}$ be an L -component algebraically connected link with admissible indexing of its components. Then for a simple Lie group G there exist positive numbers $\underline{c} = (c_1, \dots, c_L)$ and positive integers $\underline{n} = (n_1, \dots, n_{L-1})$ such that if $\underline{x} = (x_1, \dots, x_L)$ is a list of L elements of $\mathfrak{h}$ such that for any positive root $\lambda \in \Delta_+$ we have $|(\lambda, \underline{x})| \in \mathcal{A}_{\underline{c}, \underline{n}}$, then any flat connection A in the link complement $S^3 \setminus \mathcal{L}$ which satisfies the conditions*

$$f_A(\text{mer}(\mathcal{L}_j)) \sim \exp(x_j) \quad (\text{A.21})$$

is H -reducible.

Proof. We prove the theorem by induction in the number of components of $\mathcal{L}$.

Let $L = 1$. The moduli space of flat G -connections is an algebraic variety (which comes, *e.g.* from solving the Wirtinger equations (A.11) for the images of s_i under the homomorphism f_A . Therefore if the claim of the theorem does not hold, then this means that the trivial connection can be infinitesimally deformed into a non- H -reducible connection. However, this contradicts Lemma A.2 coupled with eq.(1.18).

Suppose that the theorem holds for $(L - 1)$ -component link and does not hold for an L -component link $\mathcal{L}$. Since the moduli space of flat G -connections in $S^3 \setminus \mathcal{L}$ is an algebraic variety, then this means that for any $L - 1$ positive numbers $\underline{c}_{(L)}$ and for any $L - 2$ positive integers $\underline{n}_{(L)}$ there exist L Cartan subalgebra elements $\underline{x}$ such that $x_L = 0$ and for any positive root λ we have $|(\lambda, \underline{x}_{(l)})| \in \mathcal{A}_{\underline{c}_{(L)}, \underline{n}_{(L)}}$ and there exists a flat connection A in $S^3 \setminus \mathcal{L}$ which satisfies (A.21) and is either not H -reducible, or it is H -reducible, but it can be infinitesimally deformed into a connection which is not H -reducible. Since the theorem holds for $\mathcal{L}_{(L)}$, then there exist $\underline{c}_{(L)}$ and $\underline{n}_{(L)}$ for which only the first possibility is impossible. According to Lemma A.1, there also exist such $\underline{c}_{(L)}$ and $\underline{n}_{(L)}$ for which (A.9) holds, and so according to Lemma A.2, the second possibility is also impossible. The intersection of these two areas $\mathcal{A}_{\underline{c}_{(L)}, \underline{n}_{(L)}}$ contains another area of the same form in which the assumption of induction leads to contradiction. $\square$

Proof of Theorem 1.13. This theorem is a particular case of Theorem A.3 for $G = SU(2)$. $\square$

B The first polynomials

The calculation of the polynomials $T_{j,k}^{(\pm)}$ of eqs. (2.94), (2.95) is a relatively straightforward exercise. For example, one could first present the interesting part of (2.19) as an exponential

$$\begin{aligned} & \frac{\prod_{l=1}^n (q^{-m_2-l} - q^{-\gamma_2})(q^{m_1-n+l} - 1)}{\prod_{l=1}^n (q^l - 1)} q^{-m_2(\gamma_1-1)+m_1m_2+\frac{1}{2}n(n+1)} = \\ & = \binom{m_1}{n} (1 - q^{-\gamma_2})^n q^{-m_2\gamma_1} \exp \left[\left(m_2 + m_1m_2 + \frac{1}{2}n(n+1) \right) \log(1+h) \right. \\ & \quad \left. + \sum_{l=1}^n \log \left(\frac{(1+h)^{-m_2-l} - q^{-\gamma_2}}{1 - q^{-\gamma_2}} \right) + \log \left(\frac{(1+h)^{m_1-n+l} - 1}{m_1 - n + l} \right) + \log \left(\frac{(1+h)^l - 1}{l} \right) \right]. \end{aligned} \tag{B.1}$$

The exponent should be expanded in powers of h . Since the coefficients in this expansion have a polynomial dependence on l , the sums over l produce Bernoulli polynomials of n . After calculating the exponential of the resulting power series in h , we arrive at eq.(2.94).

Here is the list of the first polynomials $T_{j,k}^{(+)}(m_1, m_2, n)$ which were obtained in this way:

$$T_{0,0}^{(+)} = 1, \quad (\text{B.2})$$

$$T_{0,1}^{(+)} = \frac{1}{2}(m_1 n + n + 2m_1 m_2 + 2m_2), \quad (\text{B.3})$$

$$T_{1,0}^{(+)} = -\frac{1}{2}(n + 2m_2 + 1), \quad (\text{B.4})$$

$$T_{0,2}^{(+)} = \frac{1}{24} \left((3m_1^2 + 5m_1 + 2)n^2 + (12m_1^2 m_2 + m_1^2 + 24m_1 m_2 - 5m_1 + 12m_2 - 6)n \right. \\ \left. + 12(m_1^2 m_2^2 + 2m_1 m_2^2 + m_2^2 - m_1 m_2 - m_2) \right), \quad (\text{B.5})$$

$$T_{1,2}^{(+)} = -\frac{1}{12} \left((3m_1 + 1)n^2 + 3(4m_1 m_2 + m_1 + 2m_2 - 1)n \right) \\ + 2(6m_1 m_2^2 + 3m_1 m_2 + 3m_2^2 - 3m_2 - 2), \quad (\text{B.6})$$

$$T_{2,2}^{(+)} = \frac{1}{24} (3n^2 + (12m_2 + 5)n + 2(6m_2^2 + 6m_2 + 1)). \quad (\text{B.7})$$

The corresponding polynomials $T_{j,k}^{(-)}$ can be either derived in a similar way from eq.(2.95) or from the list of $T_{j,k}^{(+)}$ through the relation (2.96).

C The Symmetry Principle

We are going to show that the Symmetry Principle of R. Kirby and P. Melvin [5] follows rather easily from the Main Theorem.

Theorem C.1 (R. Kirby, P. Melvin) *Let $\mathcal{L}$ be an L -component link in S^3 . For a positive integer K and for an integer j , $1 \leq j \leq L$ consider two sets of positive integer numbers $\underline{\alpha}$, $\underline{\alpha}'$ such that $1 \leq \alpha \leq K - 1$ and*

$$\alpha'_j = \alpha_k \text{ if } k \neq j, \quad \alpha'_j = K - \alpha_j \quad (\text{C.1})$$

Then

$$J_{\underline{\alpha}'}(\mathcal{L}; q) = (-1)^{\sum_{\substack{1 \leq k \leq L \\ k \neq j}} l_{jk}(\alpha_k - 1)} J_{\underline{\alpha}}(\mathcal{L}; q). \quad (\text{C.2})$$

Proof. First, we will prove this theorem for algebraically connected links with $L \geq 2$. Let $\mathcal{L}$ be such a link. It is easy to see that we can assume without loss of generality that its indexing is admissible and $j \neq 1$.

According to eqs.(1.59) and (1.64) written for the case of the empty link $\mathcal{L}'$

$$\begin{aligned} J_{\underline{\alpha}'}^{(r)}(\mathcal{L}; q) &= \frac{1}{h} q^{\text{lk}(\mathcal{L}; \underline{\alpha}')} q^{\phi_{\text{lk}}(L)} \frac{1}{\nabla_A(\mathcal{L}; q^{\underline{a}'})} \sum_{n=0}^{\infty} \frac{P'_n(\mathcal{L}; q^{\underline{a}'})}{\nabla_A^{2n}(\mathcal{L}; q^{\underline{a}'})} h^n \\ &= \frac{1}{h} q^{\text{lk}(\mathcal{L}; \underline{\alpha}')} q^{\phi_{\text{lk}}(L)} \Phi_A(\mathcal{L}; q^{\underline{a}'}) \sum_{n=0}^{\infty} F_n(q^{\underline{a}'}) h^n, \end{aligned} \quad (\text{C.3})$$

where

$$F_n(\underline{t}) = \frac{1}{\Phi_A(\mathcal{L}; \underline{t}) \nabla_A(\mathcal{L}; \underline{t})} \frac{P'_n(\mathcal{L}; \underline{t})}{\nabla_A^{2n}(\mathcal{L}; \underline{t})}. \quad (\text{C.4})$$

Since, according to relations (1.26), (1.25) and (1.62), $F_n(\underline{t})$ are rational functions of $\underline{t}$ and since $q^{\alpha'_j} = q^{-\alpha_j}$, then obviously

$$F_n(q^{\underline{a}'}) = F_n(q^{\underline{a}''}), \quad (\text{C.5})$$

where

$$\alpha''_j = \alpha_k \text{ if } k \neq j, \quad \alpha''_j = -\alpha_j. \quad (\text{C.6})$$

At the same time, eqs.(1.17) and (1.25) indicate that

$$q^{\text{lk}(\mathcal{L}; \underline{\alpha}')} \Phi_A(\mathcal{L}; q^{\underline{a}'}) = -(-1)^{\sum_{\substack{1 \leq k \leq L \\ k \neq j}} l_{jk}(\alpha_k - 1)} q^{\text{lk}(\mathcal{L}; \underline{a}'')} \Phi_A(\mathcal{L}; q^{\underline{a}''}). \quad (\text{C.7})$$

A combination of eqs.(C.3), (C.5) and (C.7) indicates that

$$J_{\underline{\alpha}'}^{(r)}(\mathcal{L}; q) = -(-1)^{\sum_{\substack{1 \leq k \leq L \\ k \neq j}} l_{jk}(\alpha_k - 1)} J_{\underline{a}''}^{(r)}(\mathcal{L}; q). \quad (\text{C.8})$$

Let us write eq.(1.68) for the case of an empty link $\mathcal{L}'$ and for the colors $\underline{\alpha}'$

$$\left[\sum_{\substack{\mu_j = \pm 1 \\ 2 \leq j \leq L \\ \mu_1 = 1}} \{ \underline{\mu} \} J_{\underline{\alpha}'}^{(r)}(\mathcal{L}; q) \right]_{(\underline{\alpha}')} = J_{\underline{\alpha}'}(\mathcal{L}; q). \quad (\text{C.9})$$

Assume that in this formula $\underline{\alpha}'$ are fixed positive numbers as described by eq.(C.1) (so that expansion in the *l.h.s.* goes in fact over the powers of h as described by eqs.(1.66), (1.67)). Then we can substitute the expression (C.8) for $J_{\underline{\alpha}'}(\mathcal{L}; q)$ in the *l.h.s.* of eq.(C.9). If we also substitute there $-\mu_j$ instead of μ_j so that $\underline{\mu}\underline{a}''$ becomes $\underline{\mu}\underline{a}$ and compare the result with eq.(C.9) written for the colors $\underline{\alpha}$, then we arrive at the relation (C.2).

Thus we proved eq.(C.2) for algebraically connected links with $L \geq 2$. For an arbitrary non-empty link $\mathcal{L}$ one can always find a knot $\mathcal{L}_0$ such that the link $\mathcal{L}_0 \cup \mathcal{L}$ is algebraically connected and has at least two components. Then eq.(C.2) holds for $\mathcal{L}_0$. If we set there $\alpha_0 = 1$, then we see that it also holds for $\mathcal{L}$, because

$$J_{1, \underline{\alpha}}(\mathcal{L}_0 \cup \mathcal{L}; q) = J_{\underline{\alpha}}(\mathcal{L}; q). \quad (\text{C.10})$$

This completes the proof of the Symmetry Principle. □

References

- [1] D. Bar-Natan, S. Garoufalidis, *On the Melvin-Morton-Rozansky conjecture*, Invent. Math. **125** (1996) 103-133.
- [2] G. Burde, H. Zieschang, *Knots*
- [3] S. Elitzur, G. Moore, A. Schwimmer, N. Seiberg, *Remarks on the canonical quantization of the Chern-Simons-Witten theory*, Nucl. Phys. **B326** (1989) 108-134.
- [4] R. Lawrence, *Asymptotic expansions of Witten-Reshetikhin-Turaev invariants for some simple 3-manifolds*, Journ. Math. Phys. **36** (1995) 6106-6129.
- [5] R. Kirby, P. Melvin, *The 3-manifold invariants of Witten and Reshetikhin-Turaev for $sl(2, \mathbb{C})$* , Invent. Math. **105** (1991) 473-545.
- [6] P. Melvin, H. Morton, *The coloured Jones function*, Commun. Math. Phys. **169** (1995) 501-520.

- [7] L. Rozansky, *On p -adic convergence of perturbative invariants of some rational homology spheres*, Duke Math. Journ. **91** (1998) 353-380.
- [8] L. Rozansky, *The universal R -matrix, Burau representation and the Melvin-Morton expansion of the colored Jones polynomial*, Adv. in Math. **134** (1998) 1-31.
- [9] V. Turaev, *Reidemeister torsion in knot theory*, Russian Math. Surveys **41** (1986) 119-182.
- [10] E. Witten, *Quantum field theory and the Jones polynomial*, Commun. Math. Phys. **121** (1989) 351-399.